

The SCHOOL of nature

IV. Learning from viruses

Alexander B. Sigalov

SignaBlok, Inc.; Shrewsbury, MA USA

Key words: multichain immune recognition receptors, MIRR, signal transduction, signaling chain homooligomerization model, SCHOOL model, cell activation, immune disorders, immune response, host-virus interactions, pattern recognition receptors, T-cell receptor, natural killer cell, natural cytotoxicity receptor, transmembrane interactions, cytoplasmic homointeractions, therapeutic targets, toll-like receptors, inhibitors, novel immunomodulators, viruses, viral pathogenesis, human cytomegalovirus, human immunodeficiency virus, HIV, human T-cell lymphotropic virus, viral entry, viral replication, viral modulation, antiviral therapy

During the co-evolution of viruses and their hosts, the latter have equipped themselves with an elaborate immune system to defend themselves from the invading viruses. In order to establish a successful infection, replicate and persist in the host, viruses have evolved numerous strategies to counter and evade host antiviral immune responses as well as exploit them for productive viral replication. These strategies include those that modulate signaling mediated by cell surface receptors. Despite tremendous advancement in recent years, the exact molecular mechanisms underlying these critical points in viral pathogenesis remain unknown. In this work, based on a novel platform of receptor signaling, the Signaling Chain HOmoOLigomerization (SCHOOL) platform, I suggest specific mechanisms used by different viruses such as human immunodeficiency virus (HIV), cytomegalovirus (CMV), severe acute respiratory syndrome coronavirus, human herpesvirus 6 and others, to modulate receptor signaling. I also use the example of HIV and CMV to illustrate how two unrelated enveloped viruses use a similar SCHOOL mechanism to modulate the host immune response mediated by two functionally different receptors: T cell antigen receptor and natural killer cell receptor, NKp30. This suggests that it is very likely that similar general mechanisms can be or are used by other viral and possibly non-viral pathogens. Learning from viruses how to target cell surface receptors not only helps us understand viral strategies to escape from the host immune surveillance, but also provides novel avenues in rational drug design and the development of new therapies for immune disorders.

Introduction

Facing the destructive consequences of microbial infections, the human immune system has evolved two arms of host defense designed to discriminate foreign agents and mount appropriate

effector responses: the innate and adaptive immune systems. Differing primarily in their receptors and receptor specificities, the innate immune system functions as the early and immediate defense mechanism and recognizes a broad set of conserved and invariant properties of non-self agents, such as viruses, through a diverse set of germ-line encoded pattern recognition receptors (PRRs), including members of the toll-like receptor (TLR) family and the retinoic acid inducible gene 1 (RIG-1)-like helicases.¹⁻³ In contrast, the adaptive arm of the immune system is the more slow-responding defense mechanism but the more pathogen-specific; infectious antigens are processed in antigen-presenting cells (APCs), presented in the context of major histocompatibility complex (MHC) class I or II molecules, and are recognized by somatically generated receptors on antigen-specific T cells that are ultimately activated and perform effector functions. Collectively, the innate and adaptive immune systems work cooperatively to defend against infection, pathogenic proliferation and disease.

In order to persist in an immunocompetent host, viruses in particular have been described to have developed intricate strategies to evade the innate immune system.⁴⁻¹¹ Following viral infection and recognition of viral components by PRRs,^{1,12-16} innate immune cells such as dendritic cells and macrophages normally respond robustly with secretion of type I interferons (IFNs), a group of pro-inflammatory cytokines that upregulate numerous interferon-stimulated genes (ISGs);^{8,17-20} overexpression of ISGs initiates a series of antiviral, antiproliferative and immunoregulatory responses against the infected cell.^{2,8,20-23} A number of viruses, including influenza and herpesvirus, employ diverse counteracting mechanisms to disrupt the IFN regulatory pathway at nearly every step, including blocking IFN induction/expression, intercepting binding of IFNs to their natural target receptors, modulating intracellular IFN-mediated signaling pathways and finally downregulation of ISG expression.²⁴⁻²⁷ By disrupting the IFN regulatory pathway, viruses are able to attenuate the antiviral properties of type I IFNs and survive recognition by the innate immune system.

Because type I IFNs also upregulate expression of MHC class I and II proteins,²⁸⁻³⁵ virus-mediated disruption of normal IFN activity has been suggested to not only interrupt innate immunity but adaptive immunity as well.² Other unrelated viruses,

*Correspondence to: Alexander B. Sigalov; Email: Sigalov@signablok.com
Submitted: 07/13/10; Revised: 08/04/10; Accepted: 08/05/10
Previously published online:
www.landesbioscience.com/journals/selfnonself/article/13279
DOI: 10.4161/self.1.4.13279




Figure 1. Targeting MIRRs: suggested immunomodulatory strategies used by viruses to enter target cells, survive and replicate. Transmembrane interactions between MIRR recognition and signaling subunits are shown by black arrows. Extracellular ligand-binding domains are shown by red. Immuno-receptor tyrosine-based activation motifs (ITAMs) are denoted by green. Circular arrow indicates viral agent-induced receptor clustering. Curved lines depict intrinsic disorder of the cytoplasmic domains of MIRR signaling subunits.

namely human immunodeficiency virus (HIV), human T-cell lymphotropic virus (HTLV) and human cytomegalovirus (CMV), have also developed strategies that modulate innate and adaptive immune processes, but do not involve type I IFNs nor IFN regulatory pathways. In contrast, HIV, HTLV and CMV target members of the family of multichain immune recognition receptors (MIRRs)³⁶⁻³⁹ found on immune cells and either disrupt or surprisingly augment MIRR-mediated activation signaling as required for self-preservation.

Functionally diverse members of the MIRR family are expressed on many different immune cells, including T and B cells, natural killer (NK) cells, mast cells, macrophages, basophils, neutrophils, eosinophils, dendritic cells (DCs) and platelets.^{36,39-41} Typical examples of MIRRs include the T-cell receptor (TCR) complex, the B-cell receptor (BCR) complex, Fc receptors (e.g., FcεRI, FcαRI, FcγRI and FcγRIII), NK receptors (e.g., NKG2D, CD94/NKG2C, KIR2DS, NKp30, NKp44 and NKp46), immunoglobulin (Ig)-like transcripts and leukocyte Ig-like receptors (ILTs and LIRs, respectively), signal regulatory

proteins (SIRPs), dendritic cell immunoactivating receptor (DCAR), myeloid DNAX adapter protein of 12 kD (DAP12)-associating lectin 1 (MDL-1), blood DC antigen 2 protein (BDCA2), novel immune-type receptor (NITR), myeloid-associated Ig-like receptor (MAIR-II), triggering receptors expressed on myeloid cells (TREM2) and the platelet collagen receptor, glycoprotein VI (GPVI). For more information on the structure and function of these and other MIRRs, I refer the reader to recent reviews.^{39,42-61} The MIRR ligand-binding subunits are integral membrane proteins with small intracellular domains that are themselves inert with regard to signaling. Signaling is achieved through the association of the ligand-binding chains with signal-transducing subunits that contain in their cytoplasmic (CYTO) domains one or more copies of the immunoreceptor tyrosine-based activation motifs (ITAMs) with two appropriately spaced tyrosines (YxxL/Ix₆₋₈YxxL/I; where x denotes non-conserved residues)⁶² or the YxxM motif,^{63,64} found in the DAP10 CYTO domain.⁶⁴ The association of the MIRR subunits in resting cells is driven mostly by the noncovalent transmembrane (TM) interactions between recognition and signaling components and plays a key role in receptor assembly, integrity and surface expression.^{40,48-50,52,57,60,65-76}

Predicted and molecularly explained by a novel model of immune signaling, the Signaling Chain HOmOligomerization (SCHOOL) model,^{37,38,40,77-84} numerous unrelated viruses employ viral proteins either to (1) disrupt intermolecular TM interactions between recognition and signaling subunits of MIRRs in an effort to disarm the receptor or (2) cluster the signaling subunits to activate or augment MIRR-triggered signaling (Fig. 1). More interesting, these viruses have exquisitely incorporated targeting and manipulation of MIRR signaling in viral processes essential to the viral life cycle: viral entry, membrane targeting and viral escape and replication. By overlapping multiple functions in a single viral protein product, the virus is able to maintain a simple genome conducive to rapid replication but have the added benefit of diverse functionality.

In this work, an intriguing principle of convergence for a number of divergent viruses in their strategic choice to uniformly target MIRRs is discussed. An investigation of how seemingly disparate viruses target a single family of membrane receptors

exposes a redundancy in viral strategies exploiting the host innate and adaptive immune systems. MIRR-targeted strategies disrupting the MIRR TM architecture from the extracellular space as well as virus-induced clustering of MIRR signaling subunits from the CYTO space (Fig. 1) are described for a select group of viruses that are functionally disparate, target different host cells and differ in their replication strategies. I will also display the power of the SCHOOL model-guided primary sequence evaluation for a number of additional viruses and its ability to predict additional MIRR-targeting viral agents not previously conceived. Furthermore, by understanding the mechanisms viruses have developed over centuries of evolution to modulate MIRR-mediated triggering in the immune response, we gain insight into the fundamental details of the mechanisms underlying normal MIRR-mediated immune activation processes and can begin to learn how to take advantage of these optimized processes. Finally, the learned viral strategies and newly developed concepts of MIRR signaling can be translated towards new lines of rational drug design efforts targeting MIRRs and modulation of immune activation.^{40,79,80,83-85} MIRR-targeted strategies stretch beyond the specific viruses discussed in this work and represent a surprising junction in viral strategies. Whether this strategy represents a convergence in evolution of disparate viruses or hints towards a similar evolutionary origin from which viruses have diverged remains to be determined.

Viruses: Classification and Pathogenesis

One of the quandaries encompassing virology and virologic discovery has been the difficulty in the classification or grouping of viruses. Although a single taxonomy governing the naming of viruses has been well-established, numerous classification methods have been suggested, highlighting similarities in virion structure, target organ systems or genomic composition. Here the principles underlying the development of the Baltimore classification method and its application towards segregating viruses based on replication methods and pathogenesis are described. However, as a consequence of viral classification and the strict segregation of viruses from one other, universal viral strategies linking differentially classified viruses have been tragically overlooked. As postulated in this report, a number of viruses that lie in different classifications are only seemingly different and generic immunomodulatory strategies targeting MIRRs serve as a surprisingly common tactic shared by them.

Viral classification. Viruses represent a collection of infectious, obligate intracellular parasites that require a living host cell to replicate. They are comprised of either DNA or RNA, a virion capsid comprised of proteins encoded by the viral nucleic acid and depending on the specific virus, a surrounding envelope. Due to the high genetic, morphologic and pathogenic variability found among different viruses, classification has proven difficult. Early attempts to organize viruses were based on their structural organization, highlighting differences in nucleic acid (DNA vs. RNA), virion symmetry, presence of an envelope and number of capsomers.^{86,87} For example, one system of viral classification^{86,87} developed by Lwoff, Horne and Tournier, the LHT

system, merged all viruses under one phylum, Vira, then divided into two subphyla, subphylum Deoxyvira (DNA viruses) and Ribovira (RNA viruses) which then divided into classes based on virion symmetry and finally segregated by the number of capsomers present in the infecting virus.

The most recent and widely accepted virus classification system is based on functional characterization that differentiates viruses based on their replication strategies and chemical nature of its nucleic acid. Coined the Baltimore classification,⁸⁸ viruses are grouped into seven groups or classes, termed the Baltimore Classes I-VII (Table 1). Each group of viruses uses a different replication strategy, such as exploitation of the host polymerases (Group I) or direct translation of injected positive-sense RNA (Group IV). Although each viral group contains viruses with the same type of nucleic acid (i.e., positive-sense single stranded (ss) RNA, double stranded (ds)DNA, etc.), there is remarkable variation in virion symmetry and presence or absence of an envelope surrounding the virus (Table 1). Therefore, viral architecture and morphology don't necessarily correlate with function and structurally different viruses unexpectedly share common functions and strategies. This section is focused on how three seemingly unrelated viruses, namely HIV, CMV and HTLV, (Table 1) share a common targeted approach in their mutual ability to modulate the immune system to enhance viral entry, replication and pathogenesis: uniform exploitation of the architecture and function of different MIRRs to directly suppress or augment immune activation (Fig. 1).

Viral pathogenesis. Despite the vast diversity in viruses and target cells in the human host, there is a common sequence of processes that serve as the foundation for all viral infection. First, the infecting virus must migrate to the primary site of infection, usually through direct inoculation or through the respiratory, gastrointestinal or genitourinary route. The virus then undergoes a process of viral entry, including attachment, a physical connection of the virus to the target cell through a viral cell recognition protein-host receptor interaction and penetration, exit from the extracellular space and entry into the cellular environment. Once inside the target cell, the virus particle uncoats and releases its viral contents, including its nucleic acid genome, in preparation of viral replication. Depending on the nature of the nucleic acid and the Baltimore group classification, viral genes may be translated directly by the host cell translation machinery (i.e., Group IV positive-sense ssRNA viruses) or incorporated into the host genome (i.e., Group I dsDNA viruses). Regardless of whether the expressing transcript originates from the viral particle itself or integrated viral genes, mRNA transcripts are translated, localize to the site of maturation and assemble into virion particles, encapsulating the viral genome in the process. Depending on the enveloped property of the infecting virus, the viral particle either surrounds itself in host membrane during budding and release (enveloped viruses) or releases without an envelope (non-enveloped viruses). Released viral progeny are then free to infect other host cells and proliferate in the host organism.

Collectively, these processes represent the fundamental stages in viral pathogenesis shared amongst members of virtually every group and class of viruses. However, inside the fine details of each

Table 1. Baltimore classification of viruses

Family	Capsid	Envelope	Genome size (kb)	Representative virus ^a	Primary target cell/organ system
Group I: dsDNA					
Adenoviridae	Icosahedral	No	26–45	Adenovirus 1	Epithelial tight junctions: heart, pancreas, nervous system, prostate, testis, lung, liver, intestine
Herpesviridae	Icosahedral	Yes	125–240	Human cytomegalovirus (CMV)	Fibroblasts
Papovaviridae	Icosahedral	No	7–8	BK Virus (polyomavirus)	Kidney epithelium, lymphocytes
Poxviridae	Ovoid	Yes	130–375	Vaccinia virus	Broad tropism
Group II: positive-sense ssDNA					
Circoviridae	Icosahedral	No	2	Transmitted transfusion virus (TTV)	Oral and intestinal mucosa
Parvoviridae	Icosahedral	No	4–6	Adeno-associated virus (AAV)	Broad tropism
Group III: dsRNA					
Bornaviridae	Icosahedral	No	5–6	Borna disease virus	Broad tropism, neuronal cells
Reoviridae	Icosahedral	No	19–32	Human rotavirus	Small intestine enterocytes
Group IV: positive-sense ssRNA					
Astroviridae	Isometric	No	6–7	Astrovirus 1	Jejunum, ileum
Calciviridae	Icosahedral	No	7–8	Norwalk virus	Upper GI tract, jejunum
Coronaviridae	Helical	Yes	28–31	SARS coronavirus (SARS-CoV)	Upper airway, alveolar epithelial cells
Flaviviridae	Spherical	Yes	10–12	Hepatitis C virus	Hepatocytes
Picornaviridae	Icosahedral	No	7–9	Hepatitis A virus	Hepatocytes, intestinal mucosa
Togaviridae	Icosahedral	Yes	10–12	Rubella virus	Nasopharynx, lymph nodes
Group V: negative-sense ssRNA					
Arenaviridae	Helical filaments	Yes	11	Lymphocytotic choriomeningitis virus (LCMV)	Broad tropism, hilar lymph nodes, lung parenchyma
				Lassa virus (LASV)	Dendritic cells, macrophages and other immune cells, hepatocytes, endothelial cells
				Mopeia virus (MOPV)	Dendritic cells, macrophages, endothelial cells
				Tacaribe virus (TACV)	Dendritic cells, macrophages
Bunyaviridae	Helical filaments	Yes	11–19	Hantaan virus	Lung parenchyma, lymph nodes, hematopoietic cells
Filoviridae	Helical filaments	Yes	19	Ebola virus	Broad tropism, mononuclear phagocytic system, mucosa
				Zaire Ebola virus (ZEBOV)	Mononuclear phagocytic system
				Sudan Ebola virus (SEBOV)	Mononuclear phagocytic system
Orthomyxoviridae	Helical filaments	Yes	10–15	Influenza virus A	Upper and lower respiratory tract
Paramyxoviridae	Helical filaments	Yes	13–18	Parainfluenza virus 1	Lower respiratory tract epithelium
Rhabdoviridae	Helical filaments	Yes	11–15	Vesicular stomatitis virus	Oral mucosa
Group VI: positive-sense ssRNA - RT					
Retroviridae	Spherical	Yes	7–13	Human immunodeficiency virus (HIV)	Intestinal mucosa, T cells, dendritic cells, macrophages, microglia
				Human T-cell lymphotropic virus type 1 (HTLV-1)	T cells
Group VII: dsDNA + RT					
Hepadnaviridae	Icosahedral	Yes	3–4	Hepatitis B virus	Hepatocytes

Abbreviations: ds, double-stranded; kb, kilobase(s); RT, reverse transcriptase; SARS, severe acute respiratory syndrome; ss, single-stranded.

^aUnderlined are the viruses discussed in this paper.

stage lay intricate subprocesses that aid in enhancing viral persistence and virulence. Unexposed until recently^{40,78-80,83-85} is the universal targeting of MIRR that multiple viruses have surreptitiously concealed in several viral processes, including viral entry, membrane targeting and viral replication. In particular, HIV and CMV specifically target different receptors within the MIRR family during viral entry through extracellular targeting mechanisms (Fig. 1A) whereas HIV and HTLV target MIRRs from the intracellular environment (Fig. 1B) during viral replication. Although these viruses selectively inhibit or augment MIRR-mediated activation of the target cell during different viral stages, viral persistence is universally enhanced either through disarmament of the immune response or enhancement of the replicative environment. By overlapping multiple processes in each viral stage, viruses have demonstrated a remarkable efficiency in their life cycle that emphasizes their advanced evolution. Intriguingly, MIRR-targeted functions enacted by viral proteins seem to be present at multiple checkpoints in viral pathogenesis, coinciding with several viral stages. Therefore, it is not unreasonable to propose that MIRRs represent a key component in the host cell that multiple viruses have ubiquitously evolved to target, disrupt or activate as desired (Fig. 1).

Viral Entry and Membrane Targeting

In order for a virus to proliferate, it must first undergo a process of attachment to the target host cell and then penetration either through fusion or direct access; collectively, these two processes comprise viral entry and are often actuated by a single protein molecule. Viral attachment has been a subject of intense investigation and several details regarding the necessary specificity of viruses for their host cells have emerged. Interestingly, disparate viruses overlap in their specificities for their primary natural receptors. For example, members of the coronaviruses (OC43),⁸⁹ orthomyxoviruses (Influenza A, B)^{90,91} and reoviruses (T3)⁹²⁻⁹⁴ contain surface receptors that are specific for sialic acid residues found on the host cell receptor whereas members of the picornaviridae (rhinoviruses, polioviruses)⁹⁵⁻⁹⁸ and retroviruses (HIV-1)⁹⁹⁻¹⁰³ bind surface receptors that adopt the canonical immunoglobulin fold such as intercellular adhesion molecule-1 (ICAM-1), the immunoglobulin G (IgG) superfamily and CD4, respectively. Although there is little sequence or structural similarity in their envelope or capsid proteins, these viruses exhibit redundancy in receptor specificity.

Following attachment, the virus penetrates the host cell either through fusion in the case of enveloped viruses or direct entry for non-enveloped viruses. Although the steps and strategies non-enveloped viruses use to enter cells are largely unknown, the events leading to viral fusion have been studied in great detail.¹⁰⁴⁻¹⁰⁹ Membrane fusion of enveloped viruses is mediated by fusion proteins that exist primarily as homo- or heterodimeric type I integral membrane proteins found embedded in the surrounding envelope.¹⁰⁸⁻¹¹⁰ Concealed in the fusion protein is the fusion sequence or fusion peptide (FP), a short hydrophobic sequence ranging from 3–6 to 24–36 amino acids, that serves as the primary mediator of virus-host cell membrane anchoring.¹⁰⁴⁻¹⁰⁶

Depending on the location of the FP and the structural nature of the fusion protein, fusion proteins are segregated in three types. Type I fusion proteins found in such viruses as influenza are comprised of alpha-helix coiled-coil domains that contain FPs at the N-terminus. Type II fusion proteins contain primarily beta-sheet structures and contain internal FP sequences. The third group of fusion proteins do not fall in the type I and II classifications and are found in such viruses as coronaviruses and herpesviruses.

After translation in the host cell, type I and II fusion proteins are fusion-incompetent and require processing by viral proteases in order to be fusion-competent or primed for fusogenic activity. Once the mature, processed and primed virus encounters a target cell, fusion events are mediated either by direct recognition and binding of the virus to its receptor on a target cell or a pH trigger commonly found in viruses that fuse within the endosome and not the outer membrane.¹⁰⁹⁻¹¹⁶ Once fusion is initiated, the fusion protein undergoes irreversible conformational changes that result in exposure of the FP. The hydrophobic peptide then embeds into the target host membrane, directly linking the virus and target cell. Previous investigation has attributed the embedding properties of the FP as a conclusion of predicted secondary sequences that FPs adopt amphipathic helices with hydrophobic residues on one face and polar residues on the opposing face.¹¹⁷ However, recent work^{79,80,83-85} has suggested that FPs from HIV and CMV not only have generalized hydrophobic sequences, but sequences that specifically target host receptors, namely members of the MIRR family. If in fact MIRR-targeted strategies are conserved in a number of viruses and overlap with viral entry, sequence analysis of FPs from viruses other than HIV and CMV should identify those viruses that share in their immunomodulatory specificities for MIRRs.

Human immunodeficiency virus. Viral entry of HIV is mediated by the product of HIV *env* expression, the type I fusion protein gp160, that is processed by HIV protease to yield the viral receptor gp120 (aa 1–511) and fusion protein gp41 (aa 512–684), found associated as heterohexameric complexes [(gp120)₃-(gp41)₃]¹¹⁸⁻¹²⁰ on the surface of HIV particles. Following encounter of a target T cell, gp120 first binds the CDR2 loop of the CD4 coreceptor. CD4 induces a conformational change in gp120 that enhances binding to a coreceptor, namely CXCR4 or CCR5, to form the ternary CD4-CXCR4/CCR5-gp120 complex.^{99,101,121-127} Consequently, membrane fusion is initiated by ternary complex-induced conformational changes in the gp120-gp41 complex that release gp41 from its metastable state and allow for the FP (aa 512–535) to integrate into the target host membrane. Once the adjoining membranes are anchored by gp41, fusion events mediated by both gp41 and gp120 occur, allowing for viral entry.

Until recently,^{40,78,85,128} the function attributed to gp41 and namely the FP has been limited to anchoring of the infecting HIV particle to the target T cell. However, it is becoming increasingly evident that the FP contributes much more to the viral pathogenesis than simply viral entry. Investigation of the primary sequence of HIV FP yields the presence of two positively charged arginines (Fig. 2). Interestingly, the TM domain (TMD) of the T cell receptor alpha chain (TCR α) also contains two positively charged residues (R, K) that lie roughly on the

Similarities in the charge distribution patterns of different immunomodulatory viral sequences




Figure 2. Primary sequence analysis of proven and predicted immunomodulatory sequences of viral fusion protein regions and other domains. Similarities in charge distribution pattern with two essential positively charged residues (shown in bold blue) spaced apart by 3–4 or 7–8 amino acids suggest a similarity of mechanisms used by diverse viruses in their pathogenesis to modulate the host immune response. Note: Although the three-dimensional structures of the analyzed sequences within the cell membrane are not known, it might be assumed that these sequences may adopt a helical conformation upon membrane binding. Thus, helical wheel projections are used for illustrative purposes only; the suggested mode of action does not depend on a particular secondary structure of the sequences. As an ideal alpha helix consists of 3.6 residues per complete turn, the angle between two residues is chosen to be 100 degrees and thus there exists a periodicity after five turns and 18 residues. For this reason, the regions shown are restricted to 18 residues. CKS-17, a synthetic retroviral envelope heptadecapeptide; FP, fusion peptide; Fr-MLV, Friend murine leukemia virus; gp, glycoprotein; HHV-6 U24, human herpesvirus 6 U24 protein; HIV, human immunodeficiency virus; HTLV-1, human T lymphotropic virus type 1; HVA, herpesvirus atelae; HVS, herpesvirus saimiri; LASV, Lassa virus; LCMV, lymphocytic choriomeningitis virus; MARV, Marburg virus; MOPV, Mopeia virus; SARS-CoV, severe acute respiratory syndrome coronavirus; SEBOV, Sudan Ebola virus; TACV, Tacaribe virus; Tip, tyrosine kinase-interacting protein; Tio, two-in-one protein; TCRα, T-cell receptor alpha chain; TMD, transmembrane domain; TMP, transmembrane peptide; ZEBOV, Zaire Ebola virus.

same face of a predicted alpha helix, being separated by 4 residues (Fig. 2). Because the TMDs of other components of the TCR, namely the CD3δ and ζ hetero- and homodimers, contain a negatively charged aspartate (D) residue, it is believed that these and other electrostatic interactions drive TCR complex formation in the largely hydrophobic environment of the TM (Fig. 3A).⁶⁶ Therefore, by having similar electrostatic properties and distribution pattern of charged residues as the TCRα TMD, HIV FP may (1) specifically bind the electronegative components of the TCR complex in a TM milieu and (2) physically and functionally disconnect the CD3δ and ζ signaling subunits from the remaining TCR complex by direct competition with the TCRα subunit (Fig. 3C).^{40,78–80,84,85} This TCR-targeted functionality of the HIV FP adds a new dimension to the binding properties of the peptide and because of the adaptive immune function associated with the TCR, compounds an immunomodulatory role.

These collective functions have been described in detail by the SCHOOL model^{37,40,78–80,84,85,129} and are becoming increasingly substantiated by emerging experimental observation.

In *in vitro* coimmunoprecipitation and fluorescence resonance energy transfer (FRET) studies, HIV FP was demonstrated to specifically associate with TCR and the gp120 ligand, CD4 and to colocalize with TCR within 50 Å.¹³⁰ Since neither gp120 nor the bulk of gp41 (aa 535–684), which contains domains thought to also interfere with T-cell activation, were included in these experiments,¹³⁰ HIV FP must contain homing sequences that drive preferential localization and binding to the TCR without any extracellular contribution; the binding specificity is limited to the TM environment and is best explained by electrostatic interactions between the HIV FP and the TCR TMDs (Fig. 3C).⁷⁸

Since gp120 is the primary HIV surface receptor that specifically binds CD4, HIV does not seemingly require gp41 or the FP particularly to serve as a binding partner for TCR or CD4. However, because the FP is heavily conserved amongst the divergent HIV subtypes, it must have other TCR-specific functions outside binding. In fact, FP was demonstrated to inhibit activation of primed lymph node cells and human T-cell lines in the presence of an activating antigen.¹³⁰ However, in the presence of phorbol 12-myristate 13-acetate (PMA)/ionomycin or mitogenic antibodies to CD3, the inhibitory activity of HIV FP was abrogated.¹³⁰ These observations of FP closely mirror those of the recently studied TCRα TM peptide (TCRα TMP or TCR TMP) and are illustrated in Figure 3B. Briefly, TCR TMP (or TCR core peptide, CP) is a 9 amino acid peptide homologous to part of the TCRα TMD and contains

the two electropositive residues (R, K) thought to be important for TCR complex formation (Fig. 2). TCR TMP was also demonstrated to have immunosuppressive effects on T cells in the presence of specific stimulating antigens,^{131–135} suggesting similar functionalities of TCR TMP and HIV FP (Fig. 3B). However, those similarities were only described in retrospective analysis of the data, leading to assignments of novel functionality to the HIV FP.

As described by the SCHOOL model,^{37,40,78–80,83–85,129} both naturally-derived HIV FP and synthetically-designed TCR TMP exploit their TM specificities for the CD3δ and ζ components of the TCR to disrupt the TM interactions that hold the TCR complex together.^{78,79,83,85} By disconnecting the recognition chains, TCRαβ, from the signaling chains, CD3δ and ζ, HIV FP functionally disrupts the TCR complex and effectively disarms the MIRR. As a consequence, when TCRαβ recognizes

and binds to its MHC-peptide partner on an APC, T-cell signaling is absent; the FP-associated signaling chains are unable to oligomerize and transduce the extracellular binding event (Fig. 3C).^{78-80,83-85}

One of the defining features of the ability of HIV to replicate and proliferate is the low fidelity of HIV reverse transcriptase (RT) that leads to high mutability and sequence variability in HIV progeny during productive infection.¹³⁶ However, the HIV gp41 FP sequence is remarkably conserved among different HIV strains, suggesting a key role of not only the need for hydrophobic residues to embed in the target membrane and permit fusion but also the two electropositive residues that mediate binding to components of the TCR. As a result, HIV FP may not only serve as a fusogenic agent, but an immunosuppressive factor targeting the TCR as well, contributing to evasion of the adaptive immune response.

Human cytomegalovirus. CMV, a member of the betaherpesvirus subfamily of herpesviruses, is an enveloped virus characterized by a large genome (196 to 241 kbp) with the capacity to encode over 160 gene products. Existing as an opportunistic pathogen, CMV proliferates during primary infection or reactivation of latent infection where an absence of effective immunity arises. Such conditions include modes where the immune system is compromised by other pathogenic agents (i.e., acquired immune deficiency syndrome, AIDS) or by prescribed immunosuppression (i.e., transplant recipients). However, the virus has also been demonstrated to replicate, reactivate and proliferate in environments where inflammation is markedly elevated.^{137,138} Although the viral factors that mediate CMV pathogenesis remain largely undetermined, three stages of CMV pathogenesis have been described: (1) stimulation of a latently infected cell to differentiate and reactivate the latent virus to replicate by proinflammatory, cytokine-driven processes, (2) immunosuppression that allows amplification of productive viral replication, either systemically or locally and (3) direct or indirect viral or host immune-mediated damage that manifests as acute or chronic disease.¹³⁹ The immunosuppression or the ability of CMV to evade and survive effector responses by innate and adaptive immune cells has been studied in great detail,¹³⁹ with novel

mechanisms targeting disruption of MIRR signaling just now emerging.

Primarily infecting fibroblasts, CMV has also been found to occupy professional APCs, namely macrophages and dendritic




Figure 3. SCHOOL insights into the mode of action of immunomodulatory viral sequences targeting T-cell receptor. (A) Structural architecture of T-cell receptor is organized by three major assembly transmembrane forces, each involving one basic and two acidic amino acid residues highlighted by blue and red, respectively (shown as a simplified axial view). (B) The experimentally observed striking similarities in characteristics and immunomodulatory activities of TCRα TMP and HIV-1 gp41 FP [Amon MA, Ali M, Bender V, Chan YN, Toth I, Manolios N. *Biochim Biophys Acta* 2006; 1763:879–88; Enk AH, Knop J. *Int Arch Allergy Immunol* 2000; 123:275–81; Wang XM, Djordjevic JT, Bender V, Manolios N. *Cell Immunol* 2002; 215:12–9; Wang XM, Djordjevic JT, Kurosaka N, Schibeci S, Lee L, Williamson P, et al. *Clin Immunol* 2002; 105:199–207; Quintana FJ, Gerber D, Kent SC, Cohen IR, Shai Y. *J Clin Invest* 2005; 115:2149–58]. (C) Within the SCHOOL model, viral agents (V) compete with the TCRα chain for binding to the CD3ε and ζ signaling subunits and thus disrupt the transmembrane interactions between the ligand-binding TCRα chain and these signaling chains. This results in disconnection and pre-dissociation of the affected signaling subunits from the remaining receptor complex and prevents formation of signaling oligomers upon multivalent antigen stimulation, thus inhibiting T-cell activation. In contrast, stimulation of this “pre-dissociated” TCR with cross-linking antibodies to signaling subunit(s) still leads to receptor triggering and cell activation. FP, fusion peptide; HIV, human immunodeficiency virus; ND, not determined; PMA, phorbol 12-myristate 13-acetate; TCRα TMP, T-cell receptor alpha chain transmembrane peptide; V, viral agent.

cells, following infection. Once inside the target host cell, CMV prepares the cell for productive replication through two mechanisms: modulation of proinflammatory IFN cytokine production and reprogramming of cellular machinery. Immediately following entry, the tegument protein pp65, stored between the virion and surrounding envelope in the mature viral particle, is released and translocates to the nucleus, reducing the level of nuclear factor kappaB (NFκB) production and blocking interferon regulatory factor-3 (IRF-3) activation.¹⁴⁰ Modulation of the IFN response is compounded by the activity of IE1-p72, a gene product expressed early after infection. By binding STAT1 and STAT2, IE1-p72 sequesters the signaling kinases and prevents their association with IRF-9, leading to the block of transcription of IFN-responsive genes.¹⁴¹ CMV also dramatically alters cellular gene expression and cell cycle progression immediately following infection, allowing for productive replication; the cell cycle is dysregulated and kept in a mitosis-like state, permitting early viral gene expression and productive replication of viral progeny before apoptosis occurs.

In addition to modulation of IFN signaling pathways in the infected cell, CMV has been described extensively to have developed mechanisms of evading the NK cell arm of the innate immune system.¹³⁹ NK cells constitute a major component of the innate immune system and are able to discriminate normal cells from those under duress or infection by monitoring the differential surface expression of MHC molecules on cells through the killer cell immunoglobulin-like receptors (KIRs). Once downregulation of MHC expression is detected, ligation of the natural cytotoxicity receptors (NCR) NKp46, NKp44 by viral hemagglutinin or NKp30 by unidentified ligands results in NK cell-mediated cytotoxicity and lysis of the affected cell. While production of MHC analogs by CMV in an infected cell to conceal the infectious process has been described in great detail,¹⁴² mechanisms of viral evasion targeting the NCR have not garnered much attention until recently.

NKp30 exists on the surface of NK cells as an NKp30-ζ receptor complex, comprised of the recognition subunit NKp30 associated with the ITAM-containing ζ signaling subunit homodimer to form a canonical MIRR. Supported by experimental evidence¹⁴³ and described by the SCHOOL model,^{40,79,80,83-85,129} ligation of the recognition subunit NKp30 and subsequent oligomerization of the ζ signaling subunit results in full activation of the MIRR. Although natural ligands for NKp30 have yet to be extensively identified, recent studies have demonstrated that the tegument protein pp65 interacts specifically and directly with the NKp30 complex, thus representing one of the first molecules to be classified as a NKp30 ligand.¹⁴³ However, rather than induce activation of the targeted NK cell, pp65 exhibits deleterious effects and inhibits NK cell activation, resulting in the inability of the NK cell to kill normal, tumor and virus-infected cells. This inhibitory effect of pp65 is explained and described to be the consequence of dissociation of the signaling ζ chains from the recognition NKp30 receptor, which renders the MIRR nonfunctional.¹⁴³ However, until recent application of the SCHOOL model,^{40,79,80,83-85} the mechanism for how binding of the NKp30-ζ complex

and dissociation of NKp30 from ζ results in the inhibition of NKp30 signaling was unknown.

Investigation of the primary sequence of the N-terminal domain of pp65 reveals the presence of several electronegative and more importantly, electropositive amino acid residues^{79,80,83-85} that may disrupt the TMD interactions between NKp30 and ζ and result in the inhibition of NK cell activation observed. By taking advantage of the presence of a negatively charged aspartate (D) in the TMD of ζ, the highly positively charged pp65 N-terminus may preferentially bind ζ through a TM interaction, effectively releasing NKp30 from its binding partner, similar to the described actions of HIV FP and TCRα TMP (Fig. 3C). Experimental evidence substantiating this mechanism will need to be demonstrated, however it is evident that CMV has developed specific mechanisms to target MIRRs redundant with other viral strategies, such as those previously described for HIV FP.^{40,78-80,84,85}

While the primary function of pp65 has been attributed to immediate inhibition of NFκB production and IRF-3 promoter-driven gene expression inside the infected cell,¹⁴⁰ pp65's effects on NK cell activity have been described as a result of extracellular exposure of pp65 to the NKp30-ζ complex¹⁴³—a quandary that needs further investigation. Whether exogenous pp65's origins come from secretion of the protein or more likely release from apoptotic cells, the membrane targeting activity of pp65 may not be as disparate from HIV FP as one would imagine, despite the non-fusogenic activity of pp65 or the major classification differences between HIV and CMV (Table 1). Demonstrated to specifically target the NKp30-ζ complex, pp65 may act identically to HIV FP in targeting an MIRR and disengaging the receptor to suppress the immune cell and permit viral persistence.

Prediction of MIRR-targeting viral agents: HTLV-1 and other viruses. Like other retroviruses, HTLV-1 enters permissive cells by binding to cellular surface molecules such as heparin sulfate proteoglycans¹⁴⁴ and the ubiquitous glucose transporter GLUT1 that serves as a receptor for both HTLV-1 and HTLV-2 viruses,¹⁴⁵⁻¹⁴⁷ followed by subsequent fusion of the viral and target cell membranes, thus releasing the viral core into the host cell cytoplasm.^{145,148-151} This fusion is mediated by several viral envelope (Env) glycoproteins that are presented on the surface of virus or infected cell as a trimer of surface (SU) glycoprotein subunits anchored to a trimer of TM glycoproteins. Remarkably, infection with cell-free HTLV-1 virions remains inefficient because naturally infected lymphocytes produce very few cell-free virions and because, of the HTLV-1 virions that are released, only 1 in 10⁵ to 10⁶ is infectious.^{146,149,150} The most efficient mode of HTLV-1 infection is cell-to-cell transmission that likely represents the sole mode of in vivo transmission for all retroviruses. Using confocal microscopy, the transfer of different HTLV-1 virion components from lymphocytes of infected patients to non-infected recipient lymphocytes has been directly visualized.¹⁵⁰

Viral fusion results from a conformational change in the TM subunit of the Env protein, triggered by the SU/receptor interaction. This engagement exposes a FP located at the N terminus of the HTLV-1 TM protein gp21.^{151,152} Similar to HIV gp41

FP, this sequence inserts into target cellular membranes and is well-known to be critical for membrane fusion activity.^{153,154} However, in contrast to the HIV FP, there has been no report to date of an immunomodulatory activity of the HTLV-1 FP.

Because T lymphocytes represent the major target cells for HTLV-1, it can be easily suggested that the TCR is a favorable target for inhibition at the viral entry stage. For these purposes, a TM-targeted strategy intended to physically and functionally disconnect TCR recognition and signaling subunits (Fig. 1A) might be effectively used by HTLV-1 as was described for HIV. The SCHOOL model^{40,79,80,83-85,129} suggests that this “secret weapon” of HTLV-1 can be represented by the viral sequence that mimics the TMD of the TCR recognition subunit (for example, the TMD of TCR α chain) and is able to insert into the cell membrane where it competes with TCR α for binding to the CD3 $\delta\epsilon$ and ζ signaling chains in the TM milieu (Fig. 3C), thereby resulting in inhibition of antigen-induced T-cell activation as with HIV FP. Through helical wheel prediction (Fig. 2) of the HTLV-1 FP, similarities in the location of electropositive residues previously described to be essential for the action of HIV FP and TCR α TMP are revealed. Positioning of the charged lysine (K) residues in HTLV-1 FP is almost identical to those for the TCR α TMP and closely resemble those of the MIRR-disrupting viral agents such as HIV FP. Therefore, it is highly likely that HTLV-1 FP targets the TCR complex in a manner identical to HIV FP and TCR α TMP and disrupts the TM interactions that hold the complex together, resulting in a defused or pre-dissociated, TCR (Fig. 3C).

Intriguingly, analysis of other seemingly unrelated viruses has yielded similar correlations in primary structure and function. Earlier studies have reported an inhibitory effect of the CKS-17 peptide on lymphocyte proliferation, a synthetic 17-mer peptide with sequence corresponding to a highly conserved region of retroviral TM proteins of human and animal retroviruses including HTLV-1.¹⁵⁵ Later, the reported immunosuppression was further confirmed and further localized to a sequence essentially identical to the sequence present in the TM protein gp21 of HTLV-1,¹⁵⁶ supporting the hypothesis that this protein participates in the mechanism of immunosuppression previously reported for the TM proteins of feline leukemia virus and other animal retroviruses.

Interestingly, peptides corresponding to regions of HIV TM protein gp41 homologous to the highly conserved and immunosuppressive sequence contained within the TM proteins p15E and gp21 of animal and human retroviruses, respectively, have been also reported to inhibit lymphoproliferation.¹⁵⁶ Recently, filoviral 17-mer peptides corresponding to a 17 amino acid domain in filoviral glycoproteins that resembles an immunosuppressive motif in retroviral envelope proteins have been demonstrated to inhibit TCR-mediated T-cell activation and cell proliferation, providing new insights in the immunopathogenesis of Ebola and Marburg viruses.¹⁵⁷ In all these peptides, a striking similarity is observed between these peptides in charged or polar residue distribution patterns with positioning of the charged lysine (K) and/or arginine (R) residues almost identical to those for the HIV FP

(Fig. 2), suggesting again a similarity in the molecular mechanisms of their immunosuppressive action.

Based on the surprising conservation in positioning of the essential electropositive residues in the helical wheel predictions of HIV-1 FP and HTLV-1 FP and its similarity to those for the TCR α TMP, it is highly probable that proteins from other unrelated viruses that also participate in viral fusion would also target MIRRs on the surface of their target cell (Fig. 1). Exploratory sequence investigation of FPs from severe acute respiratory syndrome coronavirus (SARS-CoV), Lassa virus (LASV), lymphocytic choriomeningitis virus (LCMV), Mopeia virus (MOPV) and Tacaribe virus (TACV) reveal evidence of such a hypothesis. As shown in Figure 2, there is striking similarity in the positioning of the electropositive residues on one face of the helix, despite the fact that the amino acid residues aren't necessarily conserved; for example, MOPV FP contains an arginine and lysine whereas TACV contains only lysine residues. This further supports the hypothesis of the similar molecular mechanisms suggested to underlie the immunomodulatory functions of different viruses (Fig. 1).^{80,85}

Within viral entry and immune-escape strategies (Fig. 1), intrareceptor TM interactions can be targeted not only from outside but also from inside the cell. Recently, it has been shown that the human herpesvirus 6 (HHV-6) U24 protein downregulates TCR surface expression and that U24-expressing T cells are resistant to activation by APCs.¹⁵⁸ By controlling lymphocyte activation that is often accompanied by herpesvirus reactivation, the virus might prevent its own reactivation and persist in a latent state, which is less prone to immune recognition.¹⁵⁸ In this context, U24 can serve to maintain equilibrium between the virus and its host by keeping HHV-6 titers low enough that they do not cause massive immune activation.¹⁵⁸ TCR downregulation activity has been also reported for the highly conserved membrane-proximal sequence of the tyrosine kinase-interacting protein (Tip) of herpesvirus saimiri (HVS).^{159,160} Notably, primary sequences of HHV-6 U24²⁸⁻⁶⁰ and HIV FP exhibit a similar pattern with two Arg residues spaced apart by 8 amino acids (Fig. 2). The positioning of the essential electropositive residues is remarkably conserved in HVS Tip,²¹³⁻²²⁸ the relevant domain of the two-in-one (Tio) protein of herpesvirus ateles (HVA) and HTLV-1 gp21 (Fig. 2). Thus, as recently suggested,^{80,85} the SCHOOL mechanisms similar to those applied for TCR CP and HIV gp41 FP (Fig. 3) can be used by HHV-6 and other viruses in their arsenal of immune evasion tactics. Importantly, as predicted, the viral agents prevent only antigen- but not antibody-specific T cell activation (Fig. 3C). Indeed, anti-CD3 antibodies activate HHV-6-infected T cells, resulting in great increase of viral replication.^{161,162} Interestingly, increase of viral replication induced by OKT3-mediated activation of HIV-infected T cells is currently used for purging of the latent HIV-1 reservoirs in vivo,¹⁶³ thus suggesting a potential generality of the SCHOOL mechanism-based antiviral approaches and giving the important possibility to use antibodies to MIRR signaling subunits to therapeutically modulate the affected immune cell response during viral infection.

In summary, this clearly demonstrates that viruses, despite their differences in virion structure, genomic composition or

classification, have adopted similar mechanisms (SCHOOL mechanisms) of specifically targeting MIRRs, disrupting their architecture and suppressing the immune system. Importantly, by virtue of the acquired insight into this conserved structural motif, expanded predictions, hypotheses and conclusions can be derived to begin answering the question of if shared MIRR-targeted strategies represent a conserved function or if they represent a convergent tactic of divergent viruses.

Viral Replication

Similar to viral entry, viruses have developed subprocesses targeting MIRRs that underlie other viral stages, namely viral replication, for enhancement of viral production and persistence. Following entry and uncoating in the target cell, viruses undergo an efficient and economical process of replication where copies of the viral genome are abundantly produced, viral genes are expressed and viral protein translations begin to assemble into competent viral particles. Due to the diversity in genomic structure found among the different viruses, there is also great diversity in the replication strategies they employ. Contrary to cellular genomes that are comprised uniformly of dsDNA, viral genomes span all possible structural organizations: dsDNA, dsRNA, positive-sense ssDNA, negative-sense dsDNA, positive-sense ssRNA, negative-sense ssRNA and mixed (ambisense) ssDNA or ssRNA. Consequently, viruses have developed unique replication strategies, used by the Baltimore classification method to group viruses, that require different host proteins as well as inclusion of different virally encoded proteins in their genomes. For example, group I dsDNA viruses, such as members of the adenoviral family, require host cell DNA polymerases to replicate their viral genomes and are therefore highly dependent on the replicative state of the cell; the target cell must be undergoing active replication and cell division where the cell's polymerases are most active. In contrast, group VI positive-sense ssRNA viruses, such as members of the retrovirus family, replicate their genomes by RNA-dependent DNA synthesis not by any host polymerases but by virus-encoded RT; the transcribed DNA is then used as the viral template for integration into the host genome and transcription. Because RT is not supplied by the target cell, it must be packaged with the viral progeny for further replication. Regardless of the structure and replication strategy of their genomes, all viruses express their genes as functional mRNAs early in infection and direct the cell's translational machinery to make viral proteins for eventual viral packaging.

Efficiency is essential to every viral stage but particularly to replication as it represents a pivotal point in virus production. Viruses have therefore optimized their replication strategies to exploit naturally occurring biological and cellular processes of their hosts, effectively hijacking the replication, transcription and translational machinery. However, replicative efficiency has its drawbacks; viruses are consequently dependent largely on the replicative capacity of their target cells and what functional state they are in during the infection. To overcome these limitations, several viruses have developed mechanisms of activating the infected target cell from within the CYTO environment to

enhance viral replication (Fig. 1B). This section is focused on subprocesses within the realm of viral replication that two members of the retrovirus family enact by targeting a specific MIRR, namely the TCR, from the CYTO environment. Coupled with the TM-targeted strategy (Fig. 1A) employed by FPs of HIV-1 and possibly, other viruses (Fig. 2), the CYTO-targeted strategy represents an interesting dichotomy of site of action and function that converge on the identity of the specific target.

Human immunodeficiency virus. Characterized by its positive polarity ssRNA genome and group VI classification, HIV shares a unique replicative process with other members of the retrovirus family that differs significantly from other viruses. Prior to replication, HIV virions attach to and enter T lymphocytes following formation of the ternary HIV gp120-CD4-chemokine receptor CCR4/CXCR5 complex and direct membrane fusion mediated by HIV gp41, respectively.^{99,101,121-127} Once inside the cell, the virion partially uncoats in the cytoplasm, releasing viral accessory proteins and the two copies of the positive-sense ssRNA genome housed inside the viral particle. HIV RT then initiates transcription of the viral genome, producing double-stranded cDNA transcription products that immediately associate with a number of viral (integrase, RT, matrix, Vpr)¹⁶⁴⁻¹⁶⁶ and cellular (IN1, HMGA1, BAF, EED, LEDGF/p75)¹⁶⁷ proteins to form the preintegration complex (PIC). Due to the low fidelity of HIV RT¹³⁶ that results in 3×10^{-5} mutations per replication cycle in vivo,¹⁶⁸ HIV enjoys incredible genetic diversity during virus production that closely resembles evolution but in a rapid timescale. Viral particles that introduce mutations in their genomes that exhibit increased replicative capacity will propagate and dominate the infection whereas replication-deficient variants will cease to exist.

Once formed, the PIC migrates to the nucleus by the host nuclear import machinery that only actively translocates the PIC when the cell is arrested in the G₁ phase of the cell cycle and non-dividing. Following import into the nucleus, RT-transcribed viral cDNA is integrated into the host chromosome via HIV integrase, a hallmark event that is unique to HIV. Once integrated, HIV DNA is left untranscribed in a latent stage of infection until the infected T lymphocyte is activated and coordinated interactions between HIV-encoded Tat protein, host NFκB, Sp1 transcriptional transactivating proteins and the RNA polymerase II transcriptional complex facilitate production of high levels of viral RNA.¹⁶⁹ Newly transcribed mRNAs are exported from the nucleus to the cytoplasm by HIV Rev and then translated by host ER-associated and cytoplasmic ribosomes to yield gp120 Env and Gag/Gag-Pol polyproteins, respectively. Each viral protein species translocates to the CYTO face of the plasma membrane where they associate with dimeric viral positive-sense ssRNA to form the premature viral bud that subsequently undergoes further processing, entering the final stages of viral assembly and release.

While much of the work investigating HIV replication has focused on the role of the viral regulatory protein Tat on HIV RNA transcription,¹⁷⁰⁻¹⁷³ reports have suggested a key role of cellular activating factors in enhancing replication.¹⁶⁹ In order for HIV to emerge from latent infection where the HIV genome is transcriptionally silent, the infected T lymphocyte must become

activated and initiate a signaling cascade that ultimately results in the release of NF κ B from sequestration by I κ B. Therefore, any mechanism that induces a state of activation within the infected cell would effectively enhance NF κ B activity and downstream replication of HIV. The viral accessory protein Nef has been described to affect the activation profile of CD4⁺ T lymphocytes by reducing the threshold of T-cell activation^{174,175} and also initiating a transcriptional program in Jurkat T cells similar to that of a T lymphocyte exogenously activated through the TCR.¹⁷⁶ Localization to the CYTO face of the plasma membrane seems to be required for Nef-induced activation or augmentation of activation¹⁷⁷ and association with lipid rafts and cytoplasmic signaling proteins has been proposed to play a key role.¹⁷⁸ However, details of the specific mechanisms underlying Nef-mediated augmentation of activation or reduction in threshold for activation remain largely unknown.

Originally coined “negative factor” under reports that HIV Nef reduced replication by suppressing transcription of integrated HIV genes,¹⁷⁹ it is now evident that Nef mediates several processes that collectively enhance viral replication: (1) downmodulation of surface receptors, namely CD4,^{180,181} MHC Class I proteins (HLA-A, B but not C or E),¹⁸²⁻¹⁸⁴ CD28,¹⁸⁵ and TCR in the context of simian immunodeficiency virus (SIV),¹⁸⁶ (2) enhancement of viral infectivity¹⁸⁷ and (3) modulation of signaling pathways. Among all of Nef’s functions, downmodulation of the TCR remains the most controversial and intriguing. Because of its role in initiation of the signaling cascade in T lymphocytes, TCR fills a strong potential role in Nef’s reported effects on increasing the activation state of the cell. Interestingly, HIV-2 and SIV Nef have been reported to specifically interact with the ζ signaling chain of the TCR complex but additionally induce downregulation of surface TCR from the cell surface.¹⁸⁸⁻¹⁹⁰ Functional mapping of SIV Nef has revealed that the C-terminal core domain, conserved among the different HIV-1 clades and strains, is responsible for specific ζ binding whereas the non-conserved N-terminal domain cooperatively binds AP-2 from the host thereby inducing downregulation of the bound TCR.^{189,191} Extrapolation of these results explains the lack of TCR downmodulation observed for several HIV-1 Nef variants,¹⁹² considering the genetic variability in the N-terminal domain and strengthens the observed binding data surrounding the HIV-1 Nef- ζ interaction that has been previously disputed.^{188,189}

Armed with the ability to form functional homooligomers¹⁹³⁻¹⁹⁷ on the one hand and specifically bind the signaling ζ chain of the TCR,^{188,198} on the other, HIV Nef can exert activating or augmenting effects on TCR-mediated stimulation, as described recently by the SCHOOL model.^{37,40,80,83-85,129} In contrast to extracellular targeting of the TCR by HIV FP (Fig. 1A) as described earlier in this paper, HIV Nef targets the TCR from the CYTO environment (Fig. 1B) and rather than inhibit TCR activation, enhances it. In the case of HIV FP, the signaling subunits of the TCR are physically and functionally disconnected from the recognition subunits through TMD interactions formed with HIV FP that effectively results in inhibition of antigen-mediated TCR signaling (Fig. 3C). HIV Nef may cross-link with the ζ signaling subunits through CYTO interactions

(Fig. 1B),⁴⁰ cluster TCRs and instead of disengaging the receptor, activate it or prime it for activation. While a major component of the SCHOOL model requires the ability of the TCR signaling chains to homooligomerize in receptor clusters, HIV Nef has been reported to self-oligomerize, a property already described to be vital for function.^{193-197,199,200} Therefore, through the combination of an interaction with ζ and self-oligomerization, HIV Nef may induce the formation of higher order receptor oligomers (interestingly, in intact T cells, the existence of multivalent TCR complexes responsible for sensing low concentrations of antigen has been recently reported²⁰¹) that directly activate the cell¹⁷⁶ or effectively reduce the threshold of stimulus required for full activation.^{174,175} Recent studies have indeed demonstrated clustering of HIV Nef at the immunological synapse,²⁰² the interface between the infect T lymphocyte and an APC, further supporting the notion that Nef interacts with cytoplasmic components of the TCR and likely participates in higher order oligomerization conducive to T-cell activation.

Interestingly, SIV seems to have developed additional methods of further exploiting the TCR-targeted augmentation of cellular activation Nef enacts. Characterized by rapid viral kinetics and the novel ability to replicate and proliferate in non-exogenously-stimulated macaque peripheral blood mononuclear cells (PBMC),²⁰³ SIVsmPBj, a highly pathogenic strain of SIV, induces acute, destructive disease while exhibiting an augmented replicative state.^{203,204} Underlying this disease is the presence of an ITAM sequence in SIVsmPBj Nef similar to that found in the signaling domains of the CD3 δ , CD3 ϵ , CD3 γ and ζ components of the TCR. Therefore, upon localization to the inner leaflet of the plasma membrane and association with ζ during acute infection, SIVsmPBj Nef forms high order heterooligomeric Nef- ζ complexes with significantly increased numbers of ITAM domains as compared to non-SIVsmPBj variants. Consequently, the infected cell will be prone to not only clustering of the signaling chains of the TCR by binding Nef but additional induced activation by virtue of the supplied ITAM sequences present in the viral protein. By including ITAM sequences, SIVsmPBj effectively clusters viral ITAMs with host ITAMs to induce acute activation and replication.

Despite targeting the same receptor as HIV FP, HIV Nef has the complete opposite effect on its function; rather than inactivate the receptor as observed with HIV FP, HIV Nef activates it or primes it for activation. Explained to be the result of a CYTO-targeted strategy (Fig. 1B), it is intriguing that HIV developed two mechanisms of acting on the same receptor, but eliciting different outcomes depending on the viral stage and site of action. However, through those developed viral strategies, details on how MIRRs function and initiate the intracellular cascade are revealed and provide methods of studying immune regulation but also new avenues for development of novel immunomodulatory therapeutics.

Human T cell lymphotropic virus. There is a growing line of evidence that the accessory proteins of HTLV-1 are critically involved in viral transmission and propagation and may in fact be multifunctional proteins. Key among them is the p12 protein of HTLV-1, a small oncoprotein that is produced during

the course of the natural infection *in vivo* and has been shown to have multiple functions. Analogous to the accessory HIV-1 Nef protein,^{205,206} p12 is required for optimal viral infectivity in nondividing primary lymphocytes.²⁰⁷⁻²⁰⁹ HTLV-1 viral infection of T lymphocytes is known to induce T-cell activation.²⁰⁴ As suggested, one mechanism involves activation of T cells harboring the virus and is exemplified *in vivo* by infected, non-immortalized T cell clones that display prolonged states of activation, whereas with a separate mechanism, virus-infected cells can induce activation of uninfected T cells via T-cell to T-cell interactions.²⁰⁴ In non-immortalized, HTLV-1-infected T cells, spontaneous clonal proliferation is resistant to immunosuppression by transforming growth factor β (TGF β), a cytokine implicated in terminating T-cell activation, suggesting a potential role of HTLV-1 in a defense against TGF β -induced immune suppression of the host cell.²¹⁰

Spontaneous proliferation and virus production have been reported to increase in the presence of anti-CD3 and anti-TCR antibodies while addition of HLA class I antibodies, but not HLA class II or viral proteins, shut down virus production and cell proliferation.²¹¹ These findings suggest that both virus and cell activation may occur through the TCR on the infected cell. Expression of p12 has been shown to induce nuclear factor for activation of T cells (NFAT), enhance the production of interleukin-2 (IL-2), decrease MHC-I expression, increase cytoplasmic calcium and signal transducer and activator of transcription 5 (Stat 5) activation in T cells further supporting the hypothesis that p12 may alter T-cell signaling.^{209,212-216} Interestingly, p12 is important for viral infectivity in quiescent human peripheral blood lymphocytes (PBLs) and PBMCs and the establishment of persistent infection *in vivo*, suggesting a role for p12 in the activation of quiescent lymphocytes, a prerequisite for effective viral replication *in vivo*.^{207,217} In this context, function of p12 in conditions where the majority of viral target cells are in quiescent states has been predicted to be similar to that of Nef.²⁰⁷ HTLV-1 p12-expressing cells were reported to display a decreased requirement for IL-2 to induce proliferation during suboptimal stimulation with anti-CD3 and anti-CD28 antibodies.²¹⁵ HTLV-1 replication in infected lymphocytes has also been reported to increase upon CD2 cross-linking.²¹⁸ This receptor is known to signal primarily through the associated CD3 ϵ and ζ chains.^{219,220} Studies have shown that the mitogenic activity of HTLV-1 viral particles is restricted to virus-producing T cells, requires cell-to-cell contact and may be mediated through the lymphocyte-associated antigen 3 (LFA-3)/CD2 activation pathway and that HTLV-1 virions interfere mainly with activation of peripheral T cells via CD2/ ζ but not via the CD3/TCR complex.²²¹

Overall, p12 seems to augment T-cell activation and facilitate viral replication. Thus, despite the distinct structures, both retroviral accessory proteins HTLV-1 p12 and HIV Nef are able to modulate TCR-mediated signaling and play a critical role in enhancing viral infectivity in primary lymphocytes and infected animals. Interestingly, it has been recently reported that p12 could complement for effects of Nef on HIV-1 infection of Magi-CCR5 cells, which express CD4, CXCR4 and CCR5 on the

surface or macrophages.²²² Also, Jurkat cell clones that express high levels of p12 have been found to exhibit a more rapid rate of cell proliferation than the parental cells.²²² Similarly to HIV Nef, the p12 protein, upon engagement of the TCR, localizes to the interface between T cells and antigen-presenting cells, namely the immunological synapse.²²³

Intriguingly, similarly to HIV-1 Nef protein,¹⁹⁷ HTLV-1 p12 has also been shown to form dimers.²¹⁵ It can be suggested that homooligomerization of p12 contributes to p12-mediated augmentation of T-cell activation and that molecular mechanisms of this phenomenon are similar to those that have been suggested previously for Nef through application of the SCHOOL model of TCR signaling.^{40,80} If true, the homooligomerization interface(s) of p12 represent potential therapeutic targets for antiviral treatment.

Translation of Redundant Viral Strategies into Disease Care

As depicted by members of the retroviridae and herpesviridae, namely HIV, HTLV and CMV, a wide range of viruses have developed methods of targeting members of the MIRR family of surface receptors. However, depending on the needs of the virus and at which stage of viral replication the virus is in, MIRR-induced signaling is either disrupted or enhanced. More specifically, when HIV undergoes viral entry, MIRR-triggered activation is abrogated through disruption of TM interactions in TCR by HIV FP in order to evade immune activation. Similar function is required during persistence of CMV infection where signaling through NKp30 is abrogated so as to inactivate the NK cell response and accompanying immune activation. However, where MIRR-triggered activation is needed for enhanced replication, exemplified by HIV and HTLV, viral proteins once again specifically target MIRRs, but in a concerted effort to induce triggering and subsequent cellular activation mechanisms conducive to viral production. Therefore, although viruses may be structurally different, contain different types of genomes and exhibit different replication strategies, many converge in their immune modulation strategies.

The combination of retrospective analysis of previous experiments investigating details of HIV, HTLV and CMV pathogenesis and application of a novel model of immune signaling, the SCHOOL model,^{40,79-81,83,84,129,224} has revealed a couple of key features of MIRR triggering that viruses redundantly interfere to modulate the immune response: TM interactions between the recognition and signaling subunits of MIRRs and oligomeric clustering of signaling domains. Described as TM and CYTO targets (Fig. 1), respectively, these two classes of interactions represent the foundations of MIRR triggering and provide avenues for novel but universal antiviral therapies and importantly, immunomodulatory treatment as well.

Current small molecule, antiviral research has focused on exploiting the differences between virus and host and selectively targeting a viral enzyme or process. However, due to the high mutation rate many viruses enjoy, therapies against protease or reverse transcriptase in HIV are being selected, resulting in drug

resistant viral strains that exhibit even increased pathogenicity and necessitating the discovery of novel therapeutic targets. Our discussion of the specific targeting of MIRR signaling subunits, namely the TCR ζ subunit, by HIV and HTLV provides that opportunity. Targeting of TCR-mediated signaling seems to be a shared feature of both HIV and HTLV-1 viruses and reflects a similar evolutionary pathway towards their adaptation to the host immune response that may also be shared with other unrelated viruses. Instead of inhibiting a specific enzymatic function, Nef and p12 functional targeting strategy would involve disrupting the protein-protein interface between the viral protein and the partner signaling chain to abrogate its activating function. In addition, the homointeractions between viral proteins may also emerge as a functional target since homooligomerization of viral proteins has also been shown to be essential for function. Careful investigation of the interacting surfaces on both the viral and MIRR may reveal unique features essential for binding, highlighting more rationalized drug targeting. Finally, extension of this protein-protein interaction disruption strategy should also be applied to other viruses to determine if there is increased redundancy in the processes outlined by Nef and p12. If so, MIRR-targeted antiviral research may provide a new line of generic but universal antiviral therapies.

An intriguing extension of the revealed strategies viruses redundantly use to target MIRRs is the application of them towards development of immunomodulatory agents. Viruses have evolved over thousands to millions of years and have optimized methods of disarming and evading the immune response for self-preservation. Therefore, investigation of how viruses have adapted to disarm the innate and adaptive immune system will prove invaluable in rational drug design efforts aiming to reduce immune activation or inflammation. One viral strategy, namely the disruption of TMD interactions between the signaling and recognition subunits in MIRRs suggested for HIV, HTLV, CMV and other viruses here (Fig. 1A),^{40,79,80,84,85} provides such an avenue for exquisite drug discovery and development that has the potential for rapid development. Within this approach, peptides, peptidomimetics and small molecules can be rationally designed and/or screened by using protein databases and modern technologies, including high-throughput robotic procedures, in search of effective disruptors of these specific protein-protein interactions. As such, these agents can serve as effective inhibitors of receptor signaling, thus transferring viral immune evasion strategies optimized over the millennia to therapeutic strategies that require similar functionalities.

Retrospective analysis of the primary sequence of HIV FP revealed the presence of specific electropositive residues that mirror those found in the TMD of the TCR α recognition subunit (Fig. 2).^{40,79,80,83-85} Similar findings were observed when the primary sequence of CMV pp65 was analyzed in comparison with the TMD of the NKp30 recognition subunit (data not shown). Combining those observations with functional data describing the inhibitory effect HIV FP and CMV pp65 have on TCR and NKp30 signaling, respectively, it is highly probable that they compete with the relevant recognition subunits (TCR α and

NKp30) for binding with their signaling partners. Within this context, HIV FP effectively disrupts the TCR complex and renders it useless (Fig. 3C).^{40,80,84,85} Therefore, membrane-targeted strategies mimicking those of HIV FP and CMV pp65 and exploiting the binding contribution of electropositive amino acid residues will likely have similar effects and provide useful therapies for immune disorders characterized by chronic inflammation. Coincidentally, one such avenue of TCR-targeting research has already undergone development with promising results. Derived from the primary sequence of the TCR α TMD region, synthetic hydrophobic peptides, coined the TCR TM peptides (TCR TMPs), were produced and exhibit inhibitory function in both in vitro and in vivo studies.^{131,133-135,225-230} Further studies with a D-amino acid variant also show strong efficacy, suggesting that chirality plays little role in the function of the peptide, leaving sequence pattern and electrostatics as the only mediators of function.²³¹

Although the TM-targeting strategy employed by TCR α TMP was not a prospective application based on learned viral strategies, it displays the intellectual and rational research power that can be attained by investigating what viruses and nature have already employed and optimized. Hence, we have begun to investigate the primary sequences (Fig. 2)^{78-80,84,85} of several unrelated viruses and see a remarkable homology in primary sequence and sequence pattern of a number of viral proteins, highlighting the presence of electropositive residues that may also target MIRRs. Future collaborations in bioinformatics, biochemistry and virology will undoubtedly reveal new details of the viral immune evasion strategies that are shared amongst a number of viruses that may prove useful in developing rational approaches to immune therapy.

Conclusions and Perspectives

Viral infection and the resultant immune response form a violent interplay where host homeostasis is interrupted by a propagating virus seeking to proliferate and the immune system working to quell the infection. In many cases, the virus and human host have coevolved to exist symbiotically where the virus resides in a latent phase non-pathogenic to the host. However, as new viruses emerge or crossover from other species, they will need to replicate rapidly and efficiently so as to proliferate as quickly as possible. This poses the largest pathogenic threat to humans and incurs disease that defeats the immune system and results in death of the human host. Therefore, we are forced to develop novel strategies to target the infecting virus. However, rather than targeting virus-specific proteins or processes, it would be advantageous to transfer therapeutic strategies that target redundant processes found among a number of viruses. In this work, the universal targeting of members of the MIRR family by a number of seemingly unrelated viruses that function through similar mechanisms is described. Therefore, because of recent breakthroughs in our understanding of immune signaling, it is now possible to take advantage of these general processes in drug development; the tedious work of developing virus-specific therapies would be eliminated and powerful far-reaching agents could be conceived.

In addition to the antiviral lessons learned from investigating the role of MIRRs in viral pathogenesis, several details regarding normal MIRR structure-function relationships and therapeutic intervention can be extrapolated. As demonstrated by the similar function of natural HIV FP and synthetically derived TCR α TMP, viral immune evasion strategies can be transferred to therapeutic strategies that require similar functionalities. Viruses represent years of evolution and the efficiency and optimization that come along with it. Therefore, viral functions should not only be studied as foreign processes but as efficient strategies we can use in our own attempts at immune evasion or immunomodulation.

References

1. Medzhitov R. Recognition of microorganisms and activation of the immune response. *Nature* 2007; 449:819-26.
2. Pichlmair A, Reis e Sousa C. Innate recognition of viruses. *Immunity* 2007; 27:370-83.
3. Takeuchi O, Akira S. Recognition of viruses by innate immunity. *Immunol Rev* 2007; 220:214-24.
4. Schroder M, Bowie AG. An arms race: innate antiviral responses and counteracting viral strategies. *Biochem Soc Trans* 2007; 35:1512-4.
5. Loo YM, Gale M Jr. Viral regulation and evasion of the host response. *Curr Top Microbiol Immunol* 2007; 316:295-313.
6. Keller BC, Johnson CL, Erickson AK, Gale M Jr. Innate immune evasion by hepatitis C virus and West Nile virus. *Cytokine Growth Factor Rev* 2007; 18:535-44.
7. Coscoy L. Immune evasion by Kaposi's sarcoma-associated herpesvirus. *Nat Rev Immunol* 2007; 7:391-401.
8. Takaoka A, Yanai H. Interferon signalling network in innate defence. *Cell Microbiol* 2006; 8:907-22.
9. van Wamel WJ, Rooijakkers SH, Ruyken M, van Kessel KP, van Strijp JA. The innate immune modulators staphylococcal complement inhibitor and chemotaxis inhibitory protein of *Staphylococcus aureus* are located on beta-hemolysin-converting bacteriophages. *J Bacteriol* 2006; 188:1310-5.
10. Rajagopalan S, Long EO. Viral evasion of NK-cell activation. *Trends Immunol* 2005; 26:403-5.
11. Kosugi I, Kawasaki H, Arai Y, Tsutsui Y. Innate immune responses to cytomegalovirus infection in the developing mouse brain and their evasion by virus-infected neurons. *Am J Pathol* 2002; 161:919-28.
12. Haller O, Weber F. Pathogenic viruses: smart manipulators of the interferon system. *Curr Top Microbiol Immunol* 2007; 316:315-34.
13. Saito T, Gale M Jr. Principles of intracellular viral recognition. *Curr Opin Immunol* 2007; 19:17-23.
14. Kurt-Jones EA, Popova L, Kwinn L, Haynes LM, Jones LP, Tripp RA, et al. Pattern recognition receptors TLR4 and CD14 mediate response to respiratory syncytial virus. *Nat Immunol* 2000; 1:398-401.
15. Pasare C, Medzhitov R. Toll-like receptors: linking innate and adaptive immunity. *Adv Exp Med Biol* 2005; 560:11-8.
16. Janeway CA Jr. Approaching the asymptote? Evolution and revolution in immunology. *Cold Spring Harb Symp Quant Biol* 1989; 54:1-13.
17. Doly J, Civas A, Navarro S, Uze G. Type I interferons: expression and signalization. *Cell Mol Life Sci* 1998; 54:1109-21.
18. Kunzi MS, Pitha PM. Interferon targeted genes in host defense. *Autoimmunity* 2003; 36:457-61.
19. Le Page C, Genin P, Baines MG, Hiscott J. Interferon activation and innate immunity. *Rev Immunogenet* 2000; 2:374-86.
20. Ozato K, Tailor P, Kubota T. The interferon regulatory factor family in host defense: mechanism of action. *J Biol Chem* 2007; 282:20065-9.

21. Galligan CL, Murooka TT, Rahbar R, Baig E, Majchrzak-Kita B, Fish EN. Interferons and viruses: signaling for supremacy. *Immunol Res* 2006; 35:27-40.
22. Perry AK, Chen G, Zheng D, Tang H, Cheng G. The host type I interferon response to viral and bacterial infections. *Cell Res* 2005; 15:407-22.
23. Bonjardim CA. Interferons (IFNs) are key cytokines in both innate and adaptive antiviral immune responses—and viruses counteract IFN action. *Microbes Infect* 2005; 7:569-78.
24. Cebulla CM, Miller DM, Sedmak DD. Viral inhibition of interferon signal transduction. *Intervirology* 1999; 42:325-30.
25. Garcia-Sastre A. Mechanisms of inhibition of the host interferon alpha/beta-mediated antiviral responses by viruses. *Microbes Infect* 2002; 4:647-55.
26. Goodbourn S, Didcock L, Randall RE. Interferons: cell signalling, immune modulation, antiviral response and virus countermeasures. *J Gen Virol* 2000; 81:2341-64.
27. Levy DE, Garcia-Sastre A. The virus battles: IFN induction of the antiviral state and mechanisms of viral evasion. *Cytokine Growth Factor Rev* 2001; 12:143-56.
28. Yang I, Kremen TJ, Giovannone AJ, Paik E, Odesa SK, Prins RM, et al. Modulation of major histocompatibility complex Class I molecules and major histocompatibility complex-bound immunogenic peptides induced by interferon-alpha and interferon-gamma treatment of human glioblastoma multiforme. *J Neurosurg* 2004; 100:310-9.
29. Agrawal S, Kishore MC. MHC class I gene expression and regulation. *J Hematother Stem Cell Res* 2000; 9:795-812.
30. Thomas HE, Parker JL, Schreiber RD, Kay TW. IFN-gamma action on pancreatic beta cells causes class I MHC upregulation but not diabetes. *J Clin Invest* 1998; 102:1249-57.
31. Gruschwitz MS, Vieth G. Upregulation of class II major histocompatibility complex and intercellular adhesion molecule 1 expression on scleroderma fibroblasts and endothelial cells by interferon-gamma and tumor necrosis factor alpha in the early disease stage. *Arthritis Rheum* 1997; 40:540-50.
32. Dhib-Jalbut SS, Xia Q, Drew PD, Swoveland PT. Differential upregulation of HLA class I molecules on neuronal and glial cell lines by virus infection correlates with differential induction of IFN-beta. *J Immunol* 1995; 155:2096-108.
33. Chang CH, Hammer J, Loh JE, Fodor WL, Flavell RA. The activation of major histocompatibility complex class I genes by interferon regulatory factor-1 (IRF-1). *Immunogenetics* 1992; 35:378-84.
34. Beniers AJ, Peelen WP, Debruyne FM, Schalken JA. HLA-class-I and -class-II expression on renal tumor xenografts and the relation to sensitivity for alpha-IFN, gamma-IFN and TNF. *Int J Cancer* 1991; 48:709-16.
35. Giacomini P, Fisher PB, Duigou GJ, Gambari R, Natali PG. Regulation of class II MHC gene expression by interferons: insights into the mechanism of action of interferon (review). *Anticancer Res* 1988; 8:1153-61.
36. Keegan AD, Paul WE. Multichain immune recognition receptors: similarities in structure and signaling pathways. *Immunol Today* 1992; 13:63-8.

Acknowledgements

I thank Walter Kim for his help with writing the manuscript.

Sequence Accession Numbers

Accession numbers (UniProtKB/Swiss-Prot knowledgebase, www.expasy.org/sprot) for the viruses discussed in this paper are listed below. CMV, P06725; Fr-MLV, P03390; HHV-6, Q69559; HIV-1, P04578; HTLV-1, P03381; HVA, Q9YJQ8; HVS, P22575; LASV, P08669; LCMV, P07399; MARV, P35253; MOPV, P19240; SARS-CoV, P59594; SEBOV, Q66814; TACV, P18141; ZEBOV, Q05320.

37. Sigalov A. Multi-chain immune recognition receptors: spatial organization and signal transduction. *Semin Immunol* 2005; 17:51-64.
38. Sigalov AB. Multichain immune recognition receptor signaling: different players, same game? *Trends Immunol* 2004; 25:583-9.
39. Sigalov AB, ed. Multichain Immune Recognition Receptor Signaling: From Spatiotemporal Organization to Human Disease. New York: Springer-Verlag 2008; 357.
40. Sigalov AB. Immune cell signaling: a novel mechanistic model reveals new therapeutic targets. *Trends Pharmacol Sci* 2006; 27:518-24.
41. Geijtenbeek TB, Gringhuis SI. Signalling through C-type lectin receptors: shaping immune responses. *Nat Rev Immunol* 2009; 9:465-79.
42. Krosgaard M, Davis MM. How T cells 'see' antigen. *Nat Immunol* 2005; 6:239-45.
43. DeFranco AL. B-cell activation 2000. *Immunol Rev* 2000; 176:5-9.
44. Dal Porto JM, Gauld SB, Merrell KT, Mills D, Pugh-Bernard AE, Cambier J. B cell antigen receptor signaling 101. *Mol Immunol* 2004; 41:599-613.
45. Takai T. Fc receptors and their role in immune regulation and autoimmunity. *J Clin Immunol* 2005; 25:1-18.
46. Takai T. Fc receptors: their diverse functions in immunity and immune disorders. *Springer Semin Immunopathol* 2006; 28:303-4.
47. Colonna M, Nakajima H, Navarro F, Lopez-Botet M. A novel family of Ig-like receptors for HLA class I molecules that modulate function of lymphoid and myeloid cells. *J Leukoc Biol* 1999; 66:375-81.
48. Borrego F, Kabat J, Kim DK, Lieto L, Maasho K, Pena J, et al. Structure and function of major histocompatibility complex (MHC) class I specific receptors expressed on human natural killer (NK) cells. *Mol Immunol* 2002; 38:637-60.
49. Moroi M, Jung SM. Platelet glycoprotein VI: its structure and function. *Thromb Res* 2004; 114:221-33.
50. Barclay AN, Brown MH. The SIRP family of receptors and immune regulation. *Nat Rev Immunol* 2006; 6:457-64.
51. Kanazawa N, Tashiro K, Miyachi Y. Signaling and immune regulatory role of the dendritic cell immunoreceptor (DCIR) family lectins: DCIR, DCAR, dextrin-2 and BDCA-2. *Immunobiology* 2004; 209:179-90.
52. Biassoni R, Cantoni C, Falco M, Pende D, Mollo R, Moretta L, et al. Human natural killer cell activating receptors. *Mol Immunol* 2000; 37:1015-24.
53. Biassoni R, Cantoni C, Marras D, Giron-Michel J, Falco M, Moretta L, et al. Human natural killer cell receptors: insights into their molecular function and structure. *J Cell Mol Med* 2003; 7:376-87.
54. Aoki N, Kimura S, Xing Z. Role of DAP12 in innate and adaptive immune responses. *Curr Pharm Des* 2003; 9:7-10.
55. Bakker AB, Baker E, Sutherland GR, Phillips JH, Lanier LL. Myeloid DAP12-associating lectin (MDL)-1 is a cell surface receptor involved in the activation of myeloid cells. *Proc Natl Acad Sci USA* 1999; 96:9792-6.

56. van den Berg TK, Yoder JA, Litman GW. On the origins of adaptive immunity: innate immune receptors join the tale. *Trends Immunol* 2004; 25:11-6.
57. Klesney-Tait J, Turnbull IR, Colonna M. The TREM receptor family and signal integration. *Nat Immunol* 2006; 7:1266-73.
58. Takai T. Paired immunoglobulin-like receptors and their MHC class I recognition. *Immunology* 2005; 115:433-40.
59. Nakahashi C, Tahara-Hanaoka S, Totsuka N, Okoshi Y, Takai T, Ohkohchi N, et al. Dual assemblies of an activating immune receptor, MAIR-II, with ITAM-bearing adapters DAP12 and FcRgamma chain on peritoneal macrophages. *J Immunol* 2007; 178:765-70.
60. Fujimoto M, Takatsu H, Ohno H. CMRF-35-like molecule-5 constitutes novel paired receptors, with CMRF-35-like molecule-1, to transduce activation signal upon association with FcRgamma. *Int Immunol* 2006; 18:1499-508.
61. Stewart CA, Vivier E, Colonna M. Strategies of natural killer cell recognition and signaling. *Curr Top Microbiol Immunol* 2006; 298:1-21.
62. Reth M. Antigen receptor tail clue. *Nature* 1989; 338:383-4.
63. Songyang Z, Shoelson SE, Chaudhuri M, Gish G, Pawson T, Haser WG, et al. SH2 domains recognize specific phosphopeptide sequences. *Cell* 1993; 72:767-78.
64. Wu J, Cherwinski H, Spies T, Phillips JH, Lanier LL. DAP10 and DAP12 form distinct, but functionally cooperative, receptor complexes in natural killer cells. *J Exp Med* 2000; 192:1059-68.
65. Manolios N, Bonifacio JS, Klausner RD. Transmembrane helical interactions and the assembly of the T cell receptor complex. *Science* 1990; 249:274-7.
66. Call ME, Pyrdol J, Wiedmann M, Wucherpfennig KW. The organizing principle in the formation of the T cell receptor-CD3 complex. *Cell* 2002; 111:967-79.
67. Michnoff CH, Parikh VS, Lelsz DL, Tucker PW. Mutations within the NH2-terminal transmembrane domain of membrane immunoglobulin (Ig) M alters Ig alpha and Ig beta association and signal transduction. *J Biol Chem* 1994; 269:24237-44.
68. Daron M. Fc receptor biology. *Annu Rev Immunol* 1997; 15:203-34.
69. Feng J, Garrity D, Call ME, Moffett H, Wucherpfennig KW. Convergence on a distinctive assembly mechanism by unrelated families of activating immune receptors. *Immunity* 2005; 22:427-38.
70. Feng J, Call ME, Wucherpfennig KW. The assembly of diverse immune receptors is focused on a polar membrane-embedded interaction site. *PLoS Biol* 2006; 4:142.
71. Bakema JE, de Haij S, den Hartog-Jager CF, Bakker J, Vidarsson G, van Egmond M, et al. Signaling through mutants of the IgA receptor CD89 and consequences for Fc receptor gamma-chain interaction. *J Immunol* 2006; 176:3603-10.
72. Varin-Blank N, Metzger H. Surface expression of mutated subunits of the high affinity mast cell receptor for IgE. *J Biol Chem* 1990; 265:15685-94.
73. Stevens TL, Blum JH, Foy SP, Matsuchi L, DeFranco AL. A mutation of the mu transmembrane that disrupts endoplasmic reticulum retention. Effects on association with accessory proteins and signal transduction. *J Immunol* 1994; 152:4397-406.
74. Zidovetzki R, Rost B, Pecht I. Role of transmembrane domains in the functions of B- and T-cell receptors. *Immunol Lett* 1998; 64:97-107.
75. Blum JH, Stevens TL, DeFranco AL. Role of the mu immunoglobulin heavy chain transmembrane and cytoplasmic domains in B cell antigen receptor expression and signal transduction. *J Biol Chem* 1993; 268:27236-45.
76. Ra C, Jouvin MH, Kinet JP. Complete structure of the mouse mast cell receptor for IgE (Fc epsilon RI) and surface expression of chimeric receptors (rat-mouse-human) on transfected cells. *J Biol Chem* 1989; 264:15323-7.
77. Kim WM, Sigalov AB. Viral pathogenesis, modulation of immune receptor signaling and treatment. *Adv Exp Med Biol* 2008; 640:325-49.
78. Sigalov AB. Interaction between HIV gp41 fusion peptide and T cell receptor: putting the puzzle pieces back together. *Faseb J* 2007; 21:1633-4.
79. Sigalov AB. Transmembrane interactions as immunotherapeutic targets: lessons from viral pathogenesis. *Adv Exp Med Biol* 2007; 601:335-44.
80. Sigalov AB. SCHOOL model and new targeting strategies. *Adv Exp Med Biol* 2008; 640:268-311.
81. Sigalov AB. The SCHOOL of nature. I. Transmembrane signaling. Self/Nonself—Immune Recognition and Signaling 2010; 1:4-39.
82. Sigalov AB. The SCHOOL of nature. II. Protein order, disorder and oligomerization in transmembrane signaling. Self/Nonself—Immune Recognition and Signaling 2010; 1:89-102.
83. Sigalov AB. New therapeutic strategies targeting transmembrane signal transduction in the immune system. *Cell Adh Migr* 2010; 4:255-67.
84. Sigalov AB. The SCHOOL of nature. III. From mechanistic understanding to novel therapies. Self/Nonself—Immune Recognition and Signaling 2010; 3:192-224.
85. Sigalov AB. Novel mechanistic insights into viral modulation of immune receptor signaling. *PLoS Pathog* 2009; 5:1000404.
86. Lwoff A, Tournier P. The classification of viruses. *Annu Rev Microbiol* 1966; 20:45-74.
87. Lwoff A, Horne R, Tournier P. A system of viruses. *Cold Spring Harb Symp Quant Biol* 1962; 27:51-5.
88. Baltimore D. Expression of animal virus genomes. *Bacteriol Rev* 1971; 35:235-41.
89. Vlasak R, Luytjes W, Spaan W, Palese P. Human and bovine coronaviruses recognize sialic acid-containing receptors similar to those of influenza C viruses. *Proc Natl Acad Sci USA* 1988; 85:4526-9.
90. Higa HH, Rogers GN, Paulson JC. Influenza virus hemagglutinins differentiate between receptor determinants bearing N-acetyl-, N-glycolyl- and N,O-diacetylneuraminic acids. *Virology* 1985; 144:279-82.
91. Weis W, Brown JH, Cusack S, Paulson JC, Skehel JJ, Wiley DC. Structure of the influenza virus haemagglutinin complexed with its receptor, sialic acid. *Nature* 1988; 333:426-31.
92. Gentsch JR, Pacitti AF. Differential interaction of reovirus type 3 with sialylated receptor components on animal cells. *Virology* 1987; 161:245-8.
93. Paul RW, Choi AH, Lee PW. The alpha-anomeric form of sialic acid is the minimal receptor determinant recognized by reovirus. *Virology* 1989; 172:382-5.
94. Paul RW, Lee PW. Glycophorin is the reovirus receptor on human erythrocytes. *Virology* 1987; 159:94-101.
95. Greve JM, Davis G, Meyer AM, Forte CP, Yost SC, Marlor CW, et al. The major human rhinovirus receptor is ICAM-1. *Cell* 1989; 56:839-47.
96. Mendelsohn CL, Wimmer E, Racaniello VR. Cellular receptor for poliovirus: molecular cloning, nucleotide sequence and expression of a new member of the immunoglobulin superfamily. *Cell* 1989; 56:855-65.
97. Staunton DE, Merluzzi VJ, Rothlein R, Barton R, Marlin SD, Springer TA. A cell adhesion molecule, ICAM-1, is the major surface receptor for rhinoviruses. *Cell* 1989; 56:849-53.
98. Tomassini JE, Graham D, DeWitt CM, Lineberger DW, Rodkey JA, Colonna RJ. cDNA cloning reveals that the major group rhinovirus receptor on HeLa cells is intercellular adhesion molecule 1. *Proc Natl Acad Sci USA* 1989; 86:4907-11.
99. Dalgleish AG, Beverley PC, Clapham PR, Crawford DH, Greaves MF, Weiss RA. The CD4 (T4) antigen is an essential component of the receptor for the AIDS retrovirus. *Nature* 1984; 312:763-7.
100. Klatzmann D, Champagne E, Chameret S, Gruet J, Guetard D, Hercend T, et al. T-lymphocyte T4 molecule behaves as the receptor for human retrovirus LAV. *Nature* 1984; 312:767-8.
101. Maddon PJ, Dalgleish AG, McDougal JS, Clapham PR, Weiss RA, Axel R. The T4 gene encodes the AIDS virus receptor and is expressed in the immune system and the brain. *Cell* 1986; 47:333-48.
102. McDougal JS, Kennedy MS, Sligh JM, Cort SP, Mawle A, Nicholson JK. Binding of HTLV-III/LAV to T4+ T cells by a complex of the 110K viral protein and the T4 molecule. *Science* 1986; 231:382-5.
103. Sattentau QJ, Weiss RA. The CD4 antigen: physiological ligand and HIV receptor. *Cell* 1988; 52:631-3.
104. Durell SR, Martin I, Ruyschaert JM, Shai Y, Blumenthal R. What studies of fusion peptides tell us about viral envelope glycoprotein-mediated membrane fusion (review). *Mol Membr Biol* 1997; 14:97-112.
105. Eckert DM, Kim PS. Mechanisms of viral membrane fusion and its inhibition. *Annu Rev Biochem* 2001; 70:777-810.
106. Epand RM. Fusion peptides and the mechanism of viral fusion. *Biochim Biophys Acta* 2003; 1614:116-21.
107. Teissier E, Pecheur EI. Lipids as modulators of membrane fusion mediated by viral fusion proteins. *Eur Biophys J* 2007; 36:887-99.
108. Weissenhorn W, Hinz A, Gaudin Y. Virus membrane fusion. *FEBS Lett* 2007; 581:2150-5.
109. White JM. Viral and cellular membrane fusion proteins. *Annu Rev Physiol* 1990; 52:675-97.
110. White JM. Membrane fusion. *Science* 1992; 258:917-24.
111. Daniels PS, Jeffries S, Yates P, Schild GC, Rogers GN, Paulson JC, et al. The receptor-binding and membrane-fusion properties of influenza virus variants selected using anti-haemagglutinin monoclonal antibodies. *EMBO J* 1987; 6:1459-65.
112. Hoekstra D, Kok JW. Entry mechanisms of enveloped viruses. Implications for fusion of intracellular membranes. *Biosci Rep* 1989; 9:273-305.
113. Lamb RA. Paramyxovirus fusion: a hypothesis for changes. *Virology* 1993; 197:1-11.
114. Marsh M, Helenius A. Virus entry into animal cells. *Adv Virus Res* 1989; 36:107-51.
115. Underwood PA, Skehel JJ, Wiley DC. Receptor-binding characteristics of monoclonal antibody-selected antigenic variants of influenza virus. *J Virol* 1987; 61:206-8.
116. Wiley DC, Skehel JJ. The structure and function of the hemagglutinin membrane glycoprotein of influenza virus. *Annu Rev Biochem* 1987; 56:365-94.
117. Tyler KLaF, Bernard N. Fundamental Virology. Third Edition ed. Philadelphia, PA: Lippincott—Raven Publishers 1996; 161-206.
118. Center RJ, Leapman RD, Lebowitz J, Arthur LO, Earl PL, Moss B. Oligomeric structure of the human immunodeficiency virus type 1 envelope protein on the virion surface. *J Virol* 2002; 76:7863-7.
119. Weiss CD, Levy JA, White JM. Oligomeric organization of gp120 on infectious human immunodeficiency virus type 1 particles. *J Virol* 1990; 64:5674-7.
120. Zhang CW, Chishti Y, Hussey RE, Reinherz EL. Expression, purification and characterization of recombinant HIV gp140. The gp41 ectodomain of HIV or simian immunodeficiency virus is sufficient to maintain the retroviral envelope glycoprotein as a trimer. *J Biol Chem* 2001; 276:39577-85.
121. Alkhatib G, Combadiere C, Broder CC, Feng Y, Kennedy PE, Murphy PM, et al. CC CKR5: a RANTES, MIP-1alpha, MIP-1beta receptor as a fusion cofactor for macrophage-tropic HIV-1. *Science* 1996; 272:1955-8.
122. Choe H, Farzan M, Sun Y, Sullivan N, Rollins B, Ponath PD, et al. The beta-chemokine receptors CCR3 and CCR5 facilitate infection by primary HIV-1 isolates. *Cell* 1996; 85:1135-48.
123. Deng H, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhart M, et al. Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 1996; 381:661-6.

124. Doranz BJ, Rucker J, Yi Y, Smyth RJ, Samson M, Peiper SC, et al. A dual-tropic primary HIV-1 isolate that uses fusin and the beta-chemokine receptors CKR-5, CKR-3 and CKR-2b as fusion cofactors. *Cell* 1996; 85:1149-58.
125. Feng Y, Broder CC, Kennedy PE, Berger EA. HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane, G protein-coupled receptor. *Science* 1996; 272:872-7.
126. Trkola A, Dragic T, Arthos J, Binley JM, Olson WC, Allaway GP, et al. CD4-dependent, antibody-sensitive interactions between HIV-1 and its co-receptor CCR-5. *Nature* 1996; 384:184-7.
127. Wu L, Gerard NP, Wyatt R, Choe H, Parolin C, Ruffing N, et al. CD4-induced interaction of primary HIV-1 gp120 glycoproteins with the chemokine receptor CCR-5. *Nature* 1996; 384:179-83.
128. Quintana FJ, Gerber D, Kent SC, Cohen IR, Shai Y. HIV-1 fusion peptide targets the TCR and inhibits antigen-specific T cell activation. *J Clin Invest* 2005; 115:2149-58.
129. Sigalov AB. Multichain immune recognition receptor signaling: different players, same game? *Trends Immunol* 2004; 25:583-9.
130. Quintana FJ, Gerber D, Bloch I, Cohen IR, Shai Y. A structurally altered D,L-amino acid TCRalpha transmembrane peptide interacts with the TCRalpha and inhibits T-cell activation in vitro and in an animal model. *Biochemistry* 2007; 46:2317-25.
131. Wang XM, Djordjevic JT, Kurosaka N, Schibeci S, Lee L, Williamson P, et al. T-cell antigen receptor peptides inhibit signal transduction within the membrane bilayer. *Clin Immunol* 2002; 105:199-207.
132. Wang XM, Djordjevic JT, Bender V, Manolios N. T cell antigen receptor (TCR) transmembrane peptides colocalize with TCR, not lipid rafts, in surface membranes. *Cell Immunol* 2002; 215:12-9.
133. Manolios N, Collier S, Taylor J, Pollard J, Harrison LC, Bender V. T-cell antigen receptor transmembrane peptides modulate T-cell function and T cell-mediated disease. *Nat Med* 1997; 3:84-8.
134. Amon MA, Ali M, Bender V, Chan YN, Toth I, Manolios N. Lipidation and glycosylation of a T cell antigen receptor (TCR) transmembrane hydrophobic peptide dramatically enhances in vitro and in vivo function. *Biochim Biophys Acta* 2006; 1763:879-88.
135. Manolios N, Ali M, Bender V. T-cell antigen receptor (TCR) transmembrane peptides: A new paradigm for the treatment of autoimmune diseases. *Cell Adh Migr* 2010; 4:273-83.
136. Preston BD, Poiesz BJ, Loeb LA. Fidelity of HIV-1 reverse transcriptase. *Science* 1988; 242:1168-71.
137. DeMeritt IB, Milford LE, Yurochko AD. Activation of the NFkappaB pathway in human cytomegalovirus-infected cells is necessary for efficient transactivation of the major immediate-early promoter. *J Virol* 2004; 78:4498-507.
138. Mocarski E Jr, Hahn G, White KL, et al. Myeloid cell recruitment and function in pathogenesis and latency. In: Reddehase M, Ed. *Cytomegaloviruses: Pathogenesis, Molecular Biology and Infection Control*. Norfolk, United Kingdom: Caister Scientific Press 2006; 465-82.
139. Mocarski E, Shenk T, Pass RF. Cytomegaloviruses. In: Knipe D, Howley PM, Ed. *Fields Virology* Philadelphia, PA: Lippincott Williams & Wilkins 2007; 2:2702-72.
140. Browne EP, Shenk T. Human cytomegalovirus UL83-coded pp65 virion protein inhibits antiviral gene expression in infected cells. *Proc Natl Acad Sci USA* 2003; 100:11439-44.
141. Paulus C, Krauss S, Nevels M. A human cytomegalovirus antagonist of type I IFN-dependent signal transducer and activator of transcription signaling. *Proc Natl Acad Sci USA* 2006; 103:3840-5.
142. Wiertz E, Hill A, Tortorella D, Ploegh H. Cytomegaloviruses use multiple mechanisms to elude the host immune response. *Immunol Lett* 1997; 57:213-6.
143. Arnon TI, Achdout H, Levi O, Markel G, Saleh N, Katz G, et al. Inhibition of the Nkp30 activating receptor by gp65 of human cytomegalovirus. *Nat Immunol* 2005; 6:515-23.
144. Jones KS, Fugo K, Petrow-Sadowski C, Huang Y, Bertolotto DC, Lisinski I, et al. Human T-cell leukemia virus type 1 (HTLV-1) and HTLV-2 use different receptor complexes to enter T cells. *J Virol* 2006; 80:8291-302.
145. Manel N, Taylor N, Kinet S, Kim FJ, Swainson L, Lavanya M, et al. HTLV envelopes and their receptor GLUT1, the ubiquitous glucose transporter: a new vision on HTLV infection? *Front Biosci* 2004; 9:3218-41.
146. Manel N, Kim FJ, Kinet S, Taylor N, Sitbon M, Battini JL. The ubiquitous glucose transporter GLUT-1 is a receptor for HTLV. *Cell* 2003; 115:449-59.
147. Kraft S, Kinet JP. New developments in FcepsilonRI regulation, function and inhibition. *Nat Rev Immunol* 2007; 7:365-78.
148. Pinon JD, Kelly SM, Price NC, Flanagan JU, Brighty DW. An antiviral peptide targets a coiled-coil domain of the human T-cell leukemia virus envelope glycoprotein. *J Virol* 2003; 77:3281-90.
149. Andersen PS, Geisler C, Buus S, Mariuzza RA, Karjalainen K. Role of the T cell receptor ligand affinity in T cell activation by bacterial superantigens. *J Biol Chem* 2001; 276:33452-7.
150. Igakura T, Stinchcombe JC, Goon PK, Taylor GP, Weber JN, Griffiths GM, et al. Spread of HTLV-I between lymphocytes by virus-induced polarization of the cytoskeleton. *Science* 2003; 299:1713-6.
151. Daenke S, Booth S. HTLV-1-induced cell fusion is limited at two distinct steps in the fusion pathway after receptor binding. *J Cell Sci* 2000; 113:37-44.
152. Jones PL, Korte T, Blumenthal R. Conformational changes in cell surface HIV-1 envelope glycoproteins are triggered by cooperation between cell surface CD4 and co-receptors. *J Biol Chem* 1998; 273:404-9.
153. Wilson KA, Bar S, Maerz AL, Alizon M, Pombourios P. The conserved glycine-rich segment linking the N-terminal fusion peptide to the coiled coil of human T-cell leukemia virus type 1 transmembrane glycoprotein gp21 is a determinant of membrane fusion function. *J Virol* 2005; 79:4533-9.
154. Wilson KA, Maerz AL, Pombourios P. Evidence that the transmembrane domain proximal region of the human T-cell leukemia virus type 1 fusion glycoprotein gp21 has distinct roles in the prefusion and fusion-activated states. *J Biol Chem* 2001; 276:49466-75.
155. Cianciolo GJ, Copeland TD, Oroszlan S, Snyderman R. Inhibition of lymphocyte proliferation by a synthetic peptide homologous to retroviral envelope proteins. *Science* 1985; 230:453-5.
156. Ruegg CL, Monell CR, Strand M. Inhibition of lymphoproliferation by a synthetic peptide with sequence identity to gp41 of human immunodeficiency virus type 1. *J Virol* 1989; 63:3257-60.
157. Yaddanapudi K, Palacios G, Townner JS, Chen I, Sariol CA, Nichol ST, et al. Implication of a retrovirus-like glycoprotein peptide in the immunopathogenesis of Ebola and Marburg viruses. *Faseb J* 2006; 20:2519-30.
158. Sullivan BM, Coscoy L. Downregulation of the T-cell receptor complex and impairment of T-cell activation by human herpesvirus 6 u24 protein. *J Virol* 2008; 82:602-8.
159. Min CK, Bang SY, Cho BA, Choi YH, Yang JS, Lee SH, et al. Role of amphipathic helix of a herpesviral protein in membrane deformation and T cell receptor downregulation. *PLoS Pathog* 2008; 4:000209.
160. Park J, Lee BS, Choi JK, Means RE, Choe J, Jung JU. Herpesviral protein targets a cellular WD repeat endosomal protein to downregulate T lymphocyte receptor expression. *Immunity* 2002; 17:221-33.
161. Kikuta H, Lu H, Tomizawa K, Matsumoto S. Enhancement of human herpesvirus 6 replication in adult human lymphocytes by monoclonal antibody to CD3. *J Infect Dis* 1990; 161:1085-7.
162. Roffman E, Frenkel N. Replication of human herpesvirus-6 in thymocytes activated by anti-CD3 antibody. *J Infect Dis* 1991; 164:617-8.
163. van Praag RM, Prins JM, Roos MT, Schellekens PT, Ten Berge IJ, Yong SL, et al. OKT3 and IL-2 treatment for purging of the latent HIV-1 reservoir in vivo results in selective long-lasting CD4⁺ T cell depletion. *J Clin Immunol* 2001; 21:218-26.
164. Bukrinsky MI, Sharova N, McDonald TL, Pushkarskaya T, Tarpley WG, Stevenson M. Association of integrase, matrix and reverse transcriptase antigens of human immunodeficiency virus type 1 with viral nucleic acids following acute infection. *Proc Natl Acad Sci USA* 1993; 90:6125-9.
165. Farnet CM, Haseltine WA. Determination of viral proteins present in the human immunodeficiency virus type 1 preintegration complex. *J Virol* 1991; 65:1910-5.
166. Miller MD, Farnet CM, Bushman FD. Human immunodeficiency virus type 1 preintegration complexes: studies of organization and composition. *J Virol* 1997; 71:5382-90.
167. Turlure F, Devroe E, Silver PA, Engelman A. Human cell proteins and human immunodeficiency virus DNA integration. *Front Biosci* 2004; 9:3187-208.
168. Mansky LM, Temin HM. Lower in vivo mutation rate of human immunodeficiency virus type 1 than that predicted from the fidelity of purified reverse transcriptase. *J Virol* 1995; 69:5087-94.
169. Freed EA MA. HIVs and Their Replication. In: Knipe DaH PM, Ed. *Fields Virology*. Philadelphia, PA: Lippincott Williams & Wilkins 2007; 1:12107-85.
170. Gibellini D, Vitone F, Schiavone P, Re MC. HIV-1 tat protein and cell proliferation and survival: a brief review. *New Microbiol* 2005; 28:95-109.
171. Amarapal P, Tantivanich S, Balachandra K, Matsuo K, Pitisutithum P, Chongsa-nguan M. The role of the Tat gene in the pathogenesis of HIV infection. *Southeast Asian J Trop Med Public Health* 2005; 36:352-61.
172. Seelamgari A, Maddukuri A, Berro R, de la Fuente C, Kehn K, Deng L, et al. Role of viral regulatory and accessory proteins in HIV-1 replication. *Front Biosci* 2004; 9:2388-413.
173. Strebel K. Virus-host interactions: role of HIV proteins Vif, Tat and Rev. *Aids* 2003; 17:25-34.
174. Keppler OT, Tibrioni N, Venzke S, Rauch S, Fackler OT. Modulation of specific surface receptors and activation sensitization in primary resting CD4⁺ T lymphocytes by the Nef protein of HIV-1. *J Leukoc Biol* 2006; 79:616-27.
175. Schragar JA, Marsh JW. HIV-1 Nef increases T cell activation in a stimulus-dependent manner. *Proc Natl Acad Sci USA* 1999; 96:8167-72.
176. Simmons A, Aluvihare V, McMichael A. Nef triggers a transcriptional program in T cells imitating single-signal T cell activation and inducing HIV virulence mediators. *Immunity* 2001; 14:763-77.
177. Baur AS, Sawai ET, Dazin P, Fantl WJ, Cheng-Mayer C, Peterlin BM. HIV-1 Nef leads to inhibition or activation of T cells depending on its intracellular localization. *Immunity* 1994; 1:373-84.
178. Djordjevic JT, Schibeci SD, Stewart GJ, Williamson P. HIV type 1 Nef increases the association of T cell receptor (TCR)-signaling molecules with T cell rafts and promotes activation-induced raft fusion. *AIDS Res Hum Retroviruses* 2004; 20:547-55.
179. Ahmad N, Venkatesan S. Nef protein of HIV-1 is a transcriptional repressor of HIV-1 LTR. *Science* 1988; 241:1481-5.
180. Garcia JV, Miller AD. Downregulation of cell surface CD4 by nef. *Res Virol* 1992; 143:52-5.
181. Garcia JV, Miller AD. Serine phosphorylation-independent downregulation of cell-surface CD4 by nef. *Nature* 1991; 350:508-11.
182. Schwartz O, Marechal V, Le Gall S, Lemonnier F, Heard JM. Endocytosis of major histocompatibility complex class I molecules is induced by the HIV-1 Nef protein. *Nat Med* 1996; 2:338-42.

183. Cohen GB, Gandhi RT, Davis DM, Mandelboim O, Chen BK, Strominger JL, et al. The selective down-regulation of class I major histocompatibility complex proteins by HIV-1 protects HIV-infected cells from NK cells. *Immunity* 1999; 10:661-71.
184. Le Gall S, Erdtmann L, Benichou S, Berlioz-Torrent C, Liu L, Benarous R, et al. Nef interacts with the mu subunit of clathrin adaptor complexes and reveals a cryptic sorting signal in MHC I molecules. *Immunity* 1998; 8:483-95.
185. Swigut T, Shohdy N, Skowronski J. Mechanism for downregulation of CD28 by Nef. *EMBO J* 2001; 20:1593-604.
186. Munch J, Janardhan A, Stolte N, Stahl-Hennig C, Ten Haaf P, Heeney JL, et al. T-cell receptor: CD3 downregulation is a selected in vivo function of simian immunodeficiency virus Nef but is not sufficient for effective viral replication in rhesus macaques. *J Virol* 2002; 76:12360-4.
187. Munch J, Rajan D, Schindler M, Specht A, Rucker E, Novembre FJ, et al. Nef-mediated enhancement of virion infectivity and stimulation of viral replication are fundamental properties of primate lentiviruses. *J Virol* 2007; 81:13852-64.
188. Schaefer TM, Bell I, Fallert BA, Reinhart TA. The T-cell receptor zeta chain contains two homologous domains with which simian immunodeficiency virus Nef interacts and mediates down-modulation. *J Virol* 2000; 74:3273-83.
189. Swigut T, Greenberg M, Skowronski J. Cooperative interactions of simian immunodeficiency virus Nef, AP-2 and CD3-zeta mediate the selective induction of T-cell receptor-CD3 endocytosis. *J Virol* 2003; 77:8116-26.
190. Bell I, Ashman C, Maughan J, Hooker E, Cook F, Reinhart TA. Association of simian immunodeficiency virus Nef with the T-cell receptor (TCR) zeta chain leads to TCR down-modulation. *J Gen Virol* 1998; 79:2717-27.
191. Lock M, Greenberg ME, Iafrafe AJ, Swigut T, Muench J, Kirchhoff F, et al. Two elements target SIV Nef to the AP-2 clathrin adaptor complex, but only one is required for the induction of CD4 endocytosis. *EMBO J* 1999; 18:2722-33.
192. Schindler M, Munch J, Kutsch O, Li H, Santiago ML, Bibollet-Ruche F, et al. Nef-mediated suppression of T cell activation was lost in a lentiviral lineage that gave rise to HIV-1. *Cell* 2006; 125:1055-67.
193. Ye H, Choi HJ, Poe J, Smithgall TE. Oligomerization is required for HIV-1 Nef-induced activation of the Src family protein-tyrosine kinase, Hck. *Biochemistry* 2004; 43:15775-84.
194. Hodge DR, Chen YM, Samuel KP. Oligomerization of the HIV type 2 Nef protein: mutational analysis of the heptad leucine repeat motif and cysteine residues. *AIDS Res Hum Retroviruses* 1995; 11:65-79.
195. Kienzle N, Freund J, Kalbitzer HR, Mueller-Lantzsch N. Oligomerization of the Nef protein from human immunodeficiency virus (HIV) type 1. *Eur J Biochem* 1993; 214:451-7.
196. Poe JA, Smithgall TE. HIV-1 Nef dimerization is required for Nef-mediated receptor downregulation and viral replication. *J Mol Biol* 2009; 394:329-42.
197. Arold S, Hoh F, Domergue S, Birk C, Delsuc MA, Jullien M, et al. Characterization and molecular basis of the oligomeric structure of HIV-1 nef protein. *Protein Sci* 2000; 9:1137-48.
198. Sigalov AB, Kim WM, Saline M, Stern LJ. The intrinsically disordered cytoplasmic domain of the T cell receptor zeta chain binds to the nef protein of simian immunodeficiency virus without a disorder-to-order transition. *Biochemistry* 2008; 47:12942-4.
199. Williams M, Roeth JF, Kasper MR, Filzen TM, Collins KL. Human immunodeficiency virus type 1 Nef domains required for disruption of major histocompatibility complex class I trafficking are also necessary for coprecipitation of Nef with HLA-A2. *J Virol* 2005; 79:632-6.
200. Liu LX, Heveker N, Fackler OT, Arold S, Le Gall S, Janvier K, et al. Mutation of a conserved residue (D123) required for oligomerization of human immunodeficiency virus type 1 Nef protein abolishes interaction with human thioesterase and results in impairment of Nef biological functions. *J Virol* 2000; 74:5310-9.
201. Schamel WW, Arechaga I, Risueno RM, van Santen HM, Cabezas P, Risco C, et al. Coexistence of multivalent and monovalent TCRs explains high sensitivity and wide range of response. *J Exp Med* 2005; 202:493-503.
202. Fenard D, Yonemoto W, de Noronha C, Cavois M, Williams SA, Greene WC. Nef is physically recruited into the immunological synapse and potentiates T cell activation early after TCR engagement. *J Immunol* 2005; 175:6050-7.
203. Fultz PN. Replication of an acutely lethal simian immunodeficiency virus activates and induces proliferation of lymphocytes. *J Virol* 1991; 65:4902-9.
204. Dehghani H, Brown CR, Plishka R, Buckler-White A, Hirsch VM. The ITAM in Nef influences acute pathogenesis of AIDS-inducing simian immunodeficiency viruses SIVsm and SIVagm without altering kinetics or extent of viremia. *J Virol* 2002; 76:4379-89.
205. Petit C, Buseyne F, Boccaccio C, Abastado JP, Heard JM, Schwartz O. Nef is required for efficient HIV-1 replication in cocultures of dendritic cells and lymphocytes. *Virology* 2001; 286:225-36.
206. Piguat V, Trono D. The Nef protein of primate lentiviruses. *Rev Med Virol* 1999; 9:111-20.
207. Albrecht B, Collins ND, Burniston MT, Nisbet JW, Ratner L, Green PL, et al. Human T-lymphotropic virus type 1 open reading frame I p12(I) is required for efficient viral infectivity in primary lymphocytes. *J Virol* 2000; 74:9828-35.
208. Albrecht B, D'Souza CD, Ding W, Tridandapani S, Coggeshall KM, Lairmore MD. Activation of nuclear factor of activated T cells by human T-lymphotropic virus type 1 accessory protein p12(I). *J Virol* 2002; 76:3493-501.
209. Bindhu M, Nair A, Lairmore MD. Role of accessory proteins of HTLV-1 in viral replication, T cell activation and cellular gene expression. *Front Biosci* 2004; 9:2556-76.
210. Hollsberg P, Ausubel LJ, Hafler DA. Human T cell lymphotropic virus type I-induced T cell activation. Resistance to TGFbeta1-induced suppression. *J Immunol* 1994; 153:566-73.
211. Mann DL, Martin P, Hamlin-Green G, Nalewaik R, Blattner W. Virus production and spontaneous cell proliferation in HTLV-I-infected lymphocytes. *Clin Immunol Immunopathol* 1994; 72:312-20.
212. Albrecht B, Lairmore MD. Critical role of human T-lymphotropic virus type 1 accessory proteins in viral replication and pathogenesis. *Microbiol Mol Biol Rev* 2002; 66:396-406.
213. Ding W, Albrecht B, Kelley RE, Muthusamy N, Kim SJ, Altschuld RA, et al. Human T-cell lymphotropic virus type 1 p12(I) expression increases cytoplasmic calcium to enhance the activation of nuclear factor of activated T cells. *J Virol* 2002; 76:10374-82.
214. Ding W, Kim SJ, Nair AM, Michael B, Boris-Lawrie K, Tripp A, et al. Human T-cell lymphotropic virus type 1 p12I enhances interleukin-2 production during T-cell activation. *J Virol* 2003; 77:11027-39.
215. Nicot C, Mulloy JC, Ferrari MG, Johnson JM, Fu K, Fukumoto R, et al. HTLV-1 p12(I) protein enhances STAT5 activation and decreases the interleukin-2 requirement for proliferation of primary human peripheral blood mononuclear cells. *Blood* 2001; 98:823-9.
216. Johnson JM, Mulloy JC, Ciminale V, Fullen J, Nicot C, Franchini G. The MHC class I heavy chain is a common target of the small proteins encoded by the 3' end of HTLV type 1 and HTLV type 2. *AIDS Res Hum Retroviruses* 2000; 16:1777-81.
217. Collins ND, Newbound GC, Albrecht B, Beard JL, Ratner L, Lairmore MD. Selective ablation of human T-cell lymphotropic virus type 1 p12I reduces viral infectivity in vivo. *Blood* 1998; 91:4701-7.
218. Guyot DJ, Newbound GC, Lairmore MD. Signaling via the CD2 receptor enhances HTLV-1 replication in T lymphocytes. *Virology* 1997; 234:123-9.
219. Von Bonin A, Ehrlich S, Fleischer B. The transmembrane region of CD2-associated signal-transducing proteins is crucial for the outcome of CD2-mediated T-cell activation. *Immunology* 1998; 93:376-82.
220. Wild MK, Verhagen AM, Meuer SC, Schraven B. The receptor function of CD2 in human CD2 transgenic mice is based on highly conserved associations with signal transduction molecules. *Cell Immunol* 1997; 180:168-75.
221. Kimata JT, Palker TJ, Ratner L. The mitogenic activity of human T-cell leukemia virus type I is T-cell associated and requires the CD2/LFA-3 activation pathway. *J Virol* 1993; 67:3134-41.
222. Tsukahara T, Ratner L. Substitution of HIV Type 1 Nef with HTLV-1 p12. *AIDS Res Hum Retroviruses* 2004; 20:938-43.
223. Fukumoto R, Dunder M, Nicot C, Adams A, Valeri VW, Samelson LE, et al. Inhibition of T-cell receptor signal transduction and viral expression by the linker for activation of T cells-interacting p12(I) protein of human T-cell leukemia/lymphoma virus type 1. *J Virol* 2007; 81:9088-99.
224. Sigalov AB. Signaling chain homooligomerization (SCHOOL) model. *Adv Exp Med Biol* 2008; 640:121-63.
225. Enk AH, Knop J. T cell receptor mimic peptides and their potential application in T-cell-mediated disease. *Int Arch Allergy Immunol* 2000; 123:275-81.
226. Huynh NT, Ffrench RA, Boadle RA, Manolios N. Transmembrane T-cell receptor peptides inhibit B- and natural killer-cell function. *Immunology* 2003; 108:458-64.
227. Collier S, Bolte A, Manolios N. Discrepancy in CD3-transmembrane peptide activity between in vitro and in vivo T-cell inhibition. *Scand J Immunol* 2006; 64:388-91.
228. Manolios N, Huynh NT, Collier S. Peptides in the treatment of inflammatory skin disease. *Australas J Dermatol* 2002; 43:226-7.
229. Ali M, Salam NK, Amon M, Bender V, Hibbs DE, Manolios N. T-cell antigen receptor-alpha chain transmembrane peptides: correlation between structure and function. *Int J Pept Res Ther* 2006; 12:261-7.
230. Kurosaka N, Ali M, Byth K, Manolios N. The mode of anti-arthritis peptide delivery impacts on the severity and outcome of adjuvant induced arthritis. *APLAR J Rheumatol* 2007; 10:198-203.
231. Gerber D, Quintana FJ, Bloch I, Cohen IR, Shai Y. D-enantiomer peptide of the TCRalpha transmembrane domain inhibits T-cell activation in vitro and in vivo. *Faseb J* 2005; 19:1190-2.